arXiv:1203.0493v3 [math.DG] 15 Jun 2012

SKT geometry

Gil R. Cavalcanti[*]
Department of Mathematics
Utrecht University

Abstract

SKT structures are closely related to Kähler structures, the difference being that in the Kähler case one has a complex structure which is parallel with respect to the Levi–Civita connection, while in the SKT the complex structure is parallel with respect to a metric connection with skew-symmetric and closed torsion, a concept which is a little more restrictive than asking that such connection has holonomy in $U(n)$. The inclusion of the torsion, however leaves several of the usual arguments used in Kähler geometry without a direct counterpart. We use tools from generalized complex geometry to develop the theory of SKT manifolds. We develop Hodge theory on SKT manifolds and, more generally, on parallel Hermitian manifolds and prove that their twisted cohomology inherits a $\mathbb{Z}\times\mathbb{Z}_2$-grading determined by the structure. We study Lie algebroids and differential operators associated to SKT structures and study the deformation theory of these structures. As applications we reobtain a result of Lübke and Teleman regarding the existence of SKT structures on the moduli space of instantons of a bundle over a complex surface and show that even though Kähler structures are not stable under deformations of the symplectic structure, small deformations are still SKT.

MSC classification 2010: 53C25; 53D18; 53C15.
Subject classification: Differential geometry.
Keywords: strong KT structure, generalized complex geometry, generalized Kähler geometry, Hodge theory, deformations.

Contents

1 Linear algebra **5**

2 Intrinsic torsion of generalized Hermitian structures **11**
2.1 The Nijenhuis tensor 12
2.2 The intrinsic torsion and the road to integrability 14

---

[*] `gil.cavalcanti@gmail.com`

3 SKT structures **16**

4 Parallel Hermitian and bi-Hermitian structures **21**
4.1 Parallelism and the intrinsic torsion 21

5 Hodge theory **26**
5.1 Differential operators, their adjoints and Laplacians 27
5.2 Hodge theory on SKT manifolds 28
5.3 A spectral sequence 31
5.4 Relation to Dolbeault cohomology 33
5.5 Alternative definition 36

6 Instantons over complex surfaces **40**

7 Lie algebroid differentials and other operators **43**
7.1 Lie algebroids 43
7.2 Further operators 45
7.3 Representations 50

8 Deformations **52**
8.1 First deformation problem 53
8.2 Second deformation problem and stability 54
8.3 Deformations of generalized Kähler structures 60

Introduction

Looking beyond the Levi–Civita connection in Riemannian geometry, one finds a number of other metric connections with interesting properties. Normally these families of connections are defined by some characteristic of the torsion tensor and recurrent themes of research are connections with parallel torsion or connections with skew symmetric torsion. The "strong" torsion condition refers to the last group of connections: a *strong torsion* connection on a Riemannian manifold is a metric connection whose torsion is skew symmetric and closed. In this setting, a *Kähler structure with strong torsion*, or *SKT structure* is a Hermitian structure (g, I) together with a strong torsion connection for which I is parallel. A weaker notion is that of a (strong) parallel Hermitian structure which for us means a connection with closed, skew symmetric torsion for which the holonomy is $U(n)$. The difference between a parallel Hermitian structure and an SKT structure being that for parallel Hermitian structures, integrability of the complex structure is not required.

A reason to study of such objects comes from string theory, where closed 3-forms arise naturally as fields in their sigma models [30, 31]. Once a 3-form is added to the sigma model, if one still requires a nontrivial amount of supersymmetry, the type of geometry of the target space has to move away from the usual Kähler geometry. It was precisely following this path that Gates, Hull and Roček [14] discovered the bi-Hermitian geometry that nowadays also goes by the name of generalized Kähler geometry [20] as the solutions to the $(2, 2)$-supersymmetric sigma model. Requiring less supersymmetry without giving up on the idea altogether leads one to consider models where there is more left than right supersymmetry or models where the right side is simply absent. These conditions lead to $(2,1)$ or $(2,0)$ supersymmetric sigma models and supersymmetry holds if and only if the target space has an SKT structure [23]. This point of view also leads one to consider parallel Hermitian and bi-Hermitian structures as these are geometric structures imposed by a sigma model with an extended supersymmetry algebra [9, 10].

There is yet another simple mathematical description of SKT structures: they correspond to Hermitian structures (g, I) for which the Hermitian form $\omega = g(I\cdot,\cdot)$ satisfies

$$dd^c\omega = 0,$$

where $d^c = i(\overline{\partial} - \partial)$. The 3-form $H = d^c\omega$ is the torsion of the connection mentioned earlier.

Since SKT manifolds are, in particular, complex, their space of forms inherits a natural bi-grading, but a simple check in concrete examples shows that there is no corresponding decomposition of their cohomology. Further, by studying these structures on six dimensional nilmanifolds Fino, Parton and Salamon [12] produced examples showing that their Frölicher spectral sequence does not necessarily degenerate at the second page, they do not satisfy the dd^c-lemma, manifolds carrying these structures may not be formal and that these structures are not stable under deformations of the complex structure. In short, several Kähler properties seem to have been lost once the torsion was included and nowadays much of the research in this area revolves around the search of examples of manifolds admitting SKT structures.

In fact, until now there were very few positive, conceptual results about SKT manifolds and a feature of these results was that they took the torsion 3-form into account in a substantial way. Remarkably, using the connection with closed 3-form torsion, Bismut proved a local index theorem for SKT manifolds in [2], extending previous results known for Kähler manifolds. Studying a slightly different problem, Gauduchon showed in [15] that any compact complex surface admits one SKT metric on each conformal class of Hermitian metrics, a result which proved to be fundamental in the study of non Kähler complex surfaces as carried out by Lübke and Teleman [28]. In particular, Lübke and Teleman proved that given a vector bundle over a complex Hermitian surface (with its only SKT structure), the smooth locus of the moduli space of instantons inherits an SKT structure [28] and the presence of torsion means that the metric on the moduli space is not the usual quotient metric, but something modeled on spaces transverse, but not orthogonal to the orbits. This phenomenon was re-encountered by Grantcharov, Papadopolous and Poon [18] when extending symplectic quotients to the SKT world. Indeed, there they observed that the torsion form forced them to choose two moment maps: the first cuts a submanifold, as usual, and the second guides on the choice of metric model for the quotient, which, once again, is not the orthogonal complement of the group orbits.

The last two examples were unified by the author in [7] where they were also put in the framework of reduction of Courant algebroids and generalized actions introduced in [3]. The key observation was that SKT structures have yet another description, this time, as a 'generalized structure', i.e., a geometric structure on $\mathbb{T}M = TM \oplus T^*M$. In fact, in this paper we show that, in a very precise way, SKT structures lie half way between generalized Hermitian and generalized Kähler structures. Given the success generalized geometry has had in the treatment of bihermitian structures, it is only natural to ask if it can do the same for SKT structures. In this paper we show that to be the case. In fact we show that it provides the correct point of view to study SKT structures and that several of the negative results mentioned earlier have a positive counterpart involving the torsion.

Indeed, the first observation is that, as structures defined on $\mathbb{T}M$ with the H-Courant bracket, the natural differential operator to consider is $d^H = d + H\wedge$, where $H = d^c\omega$, and ω is the Hermitian form. Hence questions about d, and its decomposition as $\partial + \overline{\partial}$ miss an important ingredient and were doomed from the start. The cohomology of d^H is only $\mathbb{Z}_2$-graded, yet we show that a parallel Hermitian structure induces a $\mathbb{Z}\times\mathbb{Z}_2$ grading on the space of forms which itself induces a $\mathbb{Z}\times\mathbb{Z}_2$ grading on the d^H-cohomology. This is achieved by introducing the *intrinsic torsion* of a generalized almost Hermitian structure and using it to show that the d^H-Laplacian preserves a $\mathbb{Z}\times\mathbb{Z}_2$-graded decomposition of the space of forms. For SKT manifolds one can go further and prove an identity of Laplacians, just as Michelsohn did for Kähler manifolds [29] and, more recently, Gualtieri, in the generalized Kähler context [19]. This way we relate a cohomology naturally defined in terms of the SKT data with the d^H-cohomology. These identities imply further that a variation of the Frölicher spectral sequence degenerates at the second page. As an application, we return to the moduli space of instantons on a bundle over a compact complex surface and show that the existence of an SKT structure in this space can be seen as a consequence of the Hodge theory developed for SKT manifolds.

Following the development of generalized complex geometry we study natural Lie algebroids associated to an SKT structure. In this case, since an SKT structure is only 'half' generalized Kähler, several of the bundles which are Lie algebroids in the generalized Kähler case cease to be in the SKT set up. Yet, one can still find several interesting operators similar to the Lie algebroid differentials but associated to subbundles of $\mathbb{T}M$ which are not involutive. These operators re-appear in our study of the problem of deformations of SKT structures. Precisely, they appear in the obstructions to deformations. Given the close proximity between SKT and generalized Kähler structures, we study the problem of deformations of generalized Kähler structures. Specifically, we show that the problem of deformations of generalized Kähler structures is obstructed and find the cohomology space where the obstructions lie. As an application of the deformation theory, we finish the paper showing that if (M,ω,I) is a compact Kähler manifold then deformations of the symplectic structure ω can always be completed with deformations of I into an generalized almost complex structure such that the pair of deformed structures is an SKT structure. In contrast, there may be no deformation of I which together with the deformed symplectic structure forms a (generalized) Kähler structure.

This paper is organized as follows. In Section 1 we develop the linear algebra pertinent to generalized complex, generalized Hermitian and SKT structures. In particular we show that an SKT structure gives rise to a $\mathbb{Z}\times\mathbb{Z}_2$-grading on the space of forms. In Section 2 we introduce the intrinsic torsion of a generalized Hermitian structure and in Sections 3 and 4 relate SKT structures and parallel Hermitian structures to the vanishing of certain components of the intrinsic torsion. In particular, we show that for an integrable SKT structure the operator d^H decomposes into a sum of three operators defined in terms of the SKT data, while for parallel Hermitian structures it decomposes as a sum of five operators. In the presence of a generalized complex extension of the SKT structure, d^H

decomposes further, into six operators. In Section 5 we study Hodge theory for SKT and parallel Hermitian structures and prove that in both cases the d^H-cohomology decomposes according to the decomposition of forms induced by the structure. As an application, in Section 6 we recover the result that the moduli space of instantons over a complex surface has an SKT structure. In Section 7 we study Lie algebroids, Lie algebroid differentials and similar operators defined on an SKT manifold and in Section 8 we use these operators to study the problem of deformations of SKT and generalized Kähler structures.

Acknowledgements: The author would like to thank Ulf Lindstrom, Martin Roček, Stefan Vandoren and Maxim Zabzine for helpful input regarding the relations to supersymmetric sigma models. This research was supported by a Marie Curie Intra European Fellowship.

1 Linear algebra

Given a vector space V^m we let $\mathbb{V} = V \oplus V^*$ be its "double". $\mathbb{V}$ is endowed with a natural symmetric pairing:

$$\langle X+\xi, Y+\eta\rangle = \frac{1}{2}(\eta(X)+\xi(Y)), \qquad X, Y \in V \ \xi, \eta \in V^*.$$

Elements of $\mathbb{V}$ act on $\wedge^\bullet V^*$ via

$$(X+\xi)\cdot\varphi = i_X\varphi + \xi\wedge\varphi.$$

One can easily check that for $v \in \mathbb{V}$

$$v\cdot(v\cdot\varphi) = \langle v, v\rangle\varphi,$$

hence $\wedge^\bullet V^*$ is naturally a module for the Clifford algebra of $\mathbb{V}$. In fact, it is the space of spinors for $Spin(\mathbb{V})$ and hence comes equipped with a spin invariant pairing, the *Chevalley pairing*:

$$(\varphi, \psi)_{Ch} = -(\varphi\wedge\psi^t)_{top},$$

where $\cdot^t$ indicates transposition, an $\mathbb{R}$-linear operator defined on decomposable forms by

$$(\alpha_1\wedge\cdots\wedge\alpha_k)^t = \alpha_k\wedge\cdots\wedge\alpha_1,$$

and *top* means taking the degree m component.

The spin group, $\mathrm{Spin}(\mathbb{V})$, acts on both $\mathbb{V}$ and on spinors in a compatible manner, namely, its action on $\mathbb{V}$ is by conjugation using Clifford multiplication

$$g_*v = gvg^{-1} \qquad \text{for all } g \in \mathrm{Spin}(\mathbb{V}), v \in \mathbb{V}$$

and on $\wedge^\bullet V^*$ by the Clifford action described above, so we have

$$g(v\cdot\varphi) = (gvg^{-1})(g\varphi) = (g_*v)(g\cdot\varphi) \qquad \text{for all } g \in \mathrm{Spin}(\mathbb{V}), v \in \mathbb{V} \text{ and } \varphi \in \wedge^\bullet V^*.$$

We will be particularly interested in the actions of 2-forms and 2-vectors, which we describe below to fix signs and notation.

Example 1.1. Given $B \in \wedge^2 V^* \subset \mathfrak{spin}(\mathbb{V})$, it acts on $\mathbb{V}$. Namely, we can regard B as a map from V into V^*

$$B(X+\xi) = -i_X B = -B(X).$$

Exponentiating this map we have

$$e_*^B : \mathbb{V} \longrightarrow \mathbb{V} \qquad e_*^B(X+\xi) = X + \xi - B(X).$$

And this action is compatible with the action of $e^B \in \mathrm{Spin}(\mathbb{V})$ on forms:

$$e^B : \varphi \longrightarrow e^B \wedge \varphi.$$

So we have $e_*^B : \wedge^k \mathbb{V} \longrightarrow \wedge^k \mathbb{V}$ and for $\alpha \in \wedge^\bullet \mathbb{V}$ and $\varphi \in \wedge^\bullet V^*$

$$(e_*^B \alpha) \cdot e^B \varphi = e^B(\alpha \cdot \varphi).$$

Example 1.2. Similarly, given $\beta \in \wedge^2 V$ we get maps

$$\beta(X+\xi) = i_\xi \beta = -\beta(\xi),$$

so for example if $\beta = \partial_{x_1} \wedge \partial_{x_2}$ and $\xi = dx_1$ then $\beta(\xi) = \partial_{x_2}$. Exponentiating, we get

$$e_*^\beta : \mathbb{V} \longrightarrow \mathbb{V} \qquad e_*^\beta(X+\xi) = X + \xi - \beta(\xi).$$

And we have a corresponding action on spinors

$$e^\beta : \wedge^\bullet V^* \longrightarrow \wedge^\bullet V^* \qquad e^\beta : \varphi \longrightarrow e^\beta \cdot \varphi,$$

where the dot indicates interior product. The convention for interior product by a multivector here is such that $\partial_{x_1} \wedge \partial_{x_2} \cdot dx_1 \wedge dx_2 = -1$.

These two actions are compatible:

$$(e_*^\beta \alpha) \cdot e^\beta \varphi = e^\beta(\alpha \cdot \varphi) \qquad \text{for all } \alpha \in \wedge^\bullet \mathbb{V}, \varphi \in \wedge^\bullet V^*. \tag{1.1}$$

We will be interested in introducing geometric structures on $\mathbb{V}$. The first we consider is a generalized metric, as introduced by Gualtieri [20].

Definition 1.3. A *generalized metric* on V is an automorphism $\mathcal{G} : \mathbb{V} \longrightarrow \mathbb{V}$ which is orthogonal and self-adjoint with respect to the natural pairing and for which the bilinear tensor

$$\langle \mathcal{G}v, w \rangle, \qquad v, w \in \mathbb{V}$$

is positive definite.

Since $\mathcal{G}$ is orthogonal and self-adjoint, we have $\mathcal{G}^{-1} = \mathcal{G}^t = \mathcal{G}$, hence $\mathcal{G}^2 = \mathrm{Id}$. Therefore $\mathcal{G}$ splits $\mathbb{V}$ into its ±1-eigenspaces: $\mathbb{V} = V_+ \oplus V_-$ and the projection $\pi_V : \mathbb{V} \longrightarrow V$ gives isomorphisms $\pi : V_\pm \longrightarrow V$. Further, given a generalized metric $\mathcal{G}$ we can write $\mathbb{V} = \mathcal{G}V^* \oplus V^*$ and $\mathcal{G}V^*$ is isomorphic to V via the projection $\pi_V : \mathbb{V} \longrightarrow V$. Since both V and $\mathcal{G}V$ are isotropic subspaces of $\mathbb{V}$ which project isomorphically onto V, we can describe $\mathcal{G}V$ as the graph of a linear map $b : V \longrightarrow V^*$, that is $b \in \otimes^2 V^*$. Isotropy means that

$b \in \wedge^2 V^*$ and hence gives rise to an orthogonal transformation of the natural pairing, e_*^b. This map has the property that $e_*^b : \mathcal{G}V^* \longrightarrow V$, hence, after an orthogonal transformation of $\mathbb{V}$, we can assume that $\mathcal{G}V^* = V$. For this splitting,

$$\mathcal{G} = \begin{pmatrix} 0 & g \\ g^{-1} & 0 \end{pmatrix}$$

where g is an ordinary metric on V. The splitting of $\mathbb{V}$ determined by a generalized metric is the *metric splitting*.

If V is endowed with an orientation, we can define a generalized *Hodge star operator* [19] as follows. Since $\pi_V : V_+ \longrightarrow V$ is an isomorphism, V_+ inherits an orientation. Then we let $\{e_1, e_2, \cdots, e_m\}$ be a positive orthonormal basis of V_+ and let $\star = -e_m \cdots e_2 \cdot e_1 \in \mathrm{Clif}(\mathbb{V})$. Then

$$\star\varphi := \star \cdot \varphi,$$

where $\cdot$ denotes Clifford action.

With this definition, we have

$$(\varphi, \star\varphi)_{Ch} > 0 \qquad \text{if } \varphi \neq 0 \tag{1.2}$$

If the splitting of $\mathbb{V}$ is the metric splitting, we have

$$(\varphi, \star\psi)_{Ch} = \varphi \wedge *\psi,$$

where $*$ is the usual Hodge star, hence, in this splitting, $\star$ is the usual Hodge star except for a change in signs given by the Chevalley pairing. Since $\star = -e_m \cdot \cdots \cdot e_2 \cdot e_1$, we have that

$$\star^2 = (-1)^{\frac{m(m-1)}{2}}$$

and hence it splits the space of forms into its eigenspaces, namely, into its ± 1-eigenspaces $\wedge^{\bullet}_{\pm} V^*$ if m is zero or one modulo four or its $\pm i$-eigenspaces if n is 2 or 3 modulo 4. This allows us to define *self-dual* and *anti self-dual forms* in dimensions zero and one modulo four. However we would like to have a single definition for all dimensions, so we introduce the concept of SD- and ASD-forms:

Definition 1.4. We say that a form $\varphi \in \wedge^{\bullet} V^*_{\mathbb{C}}$ is SD if $\star\varphi = -i^{\frac{m(m-1)}{2}} \varphi$ and that it is ASD if $\star\varphi = i^{\frac{m(m-1)}{2}} \varphi$. We denote the space of SD-forms by $\wedge^{\bullet}_{+} V^*$ and the space of ASD-forms by $\wedge^{\bullet}_{-} V^*$.[1]

The Clifford action of elements in $V_\pm$ either preserves or switches the eigenspaces of $\star$:

Lemma 1.5. Let $v_\pm \in V_\pm$. Then acting via Clifford action on forms we have

$$v_+ \star = (-1)^{m-1} \star v_+ \qquad \text{and} \qquad v_- \star = (-1)^m \star v_-$$

hence for m even,

$$V_+ : \wedge^{\bullet}_{\pm} V^* \longrightarrow \wedge^{\bullet}_{\mp} V^* \qquad \text{and} \qquad V_- : \wedge^{\bullet}_{\pm} V^* \longrightarrow \wedge^{\bullet}_{\pm} V^*$$

---

[1] The choice of signs on the Chevalley pairing and of $\star$ were made so that on a four manifold, the notion of SD and ASD agrees with the usual notions of self-dual and anti self-dual on 2-forms.

and for m odd

$$V_+ : \wedge^\bullet_\pm V^* \longrightarrow \wedge^\bullet_\pm V^* \qquad \text{and} \qquad V_- : \wedge^\bullet_\pm V^* \longrightarrow \wedge^\bullet_\mp V^*$$

Proof. For $v_- \in V_-$, we have that $\langle v_-, V_+\rangle = 0$ hence v_- graded commutes with $\star$. For $v_+ \in V_+$, we can choose an orthonormal basis for V_+ for which $e_1 = v/\|v\|$, so that v anti commutes with all the remaining elements of the basis and commutes with e_1. □

For the rest of this section we will introduce structures on V which force its dimension to be even so we let $m = 2n$.

Definition 1.6. A *generalized complex structure* on V is a complex structure on $\mathbb{V}$ which is orthogonal with respect to the natural pairing. A *generalized Hermitian structure* or a $U(n) \times U(n)$ *structure* on V is a generalized complex structure $\mathcal{J}_1$ on V and a generalized metric $\mathcal{G}$ such that $\mathcal{J}_1$ and $\mathcal{G}$ commute.

Given a generalized complex structure $\mathcal{J}$ on V, we can split $\mathbb{V}_{\mathbb{C}}$, the complexification of $\mathbb{V}$, into the $\pm i$-eigenspaces of $\mathcal{J}$: $\mathbb{V}_{\mathbb{C}} = L \oplus \overline{L}$. The spaces L and $\overline{L}$ are maximal isotropic subspaces of $\mathbb{V}_{\mathbb{C}}$ such that $L \cap \overline{L} = \{0\}$. Since the natural pairing is nondegenerate, we can use it to identify $\overline{L} = L^*$. Precisely, we identify

$$v \mapsto 2\langle v, \cdot\rangle \in L^*, \qquad \text{for } v \in \overline{L}. \tag{1.3}$$

Given a generalized Hermitian structure $(\mathcal{J}_1, \mathcal{G})$ on V, $\mathcal{J}_2 = \mathcal{G}\mathcal{J}_1$ is orthogonal with respect to the natural pairing and squares to $-\mathrm{Id}$, hence it is also a generalized complex structure. Since $\pi_V : V_\pm \longrightarrow V$ are isomorphisms, and $\mathcal{J}_1|_{V_\pm}$ is a complex structure on $V_\pm$ orthogonal with respect to the natural pairing, it induces complex structures $I_\pm$ on V compatible with the metric g induced by $\mathcal{G}$ making V into a bi-Hermitian vector space. We can further form the corresponding Hermitian forms $\omega_\pm = g \circ I_\pm$.

Given any generalized Hermitian structure, $\wedge^\bullet V^*_{\mathbb{C}}$ splits as the intersections of the eigenspaces of $\mathcal{J}_1$ and $\mathcal{J}_2$: $U^{p,q} = U^p_{\mathcal{J}_1} \cap U^q_{\mathcal{J}_2}$, where $U^p_{\mathcal{J}_i}$ is the ip-eigenspace of $\mathcal{J}_i$ in $\wedge^\bullet V^*_{\mathbb{C}}$. In this context, the generalized Hodge star is related to the action of $\mathbb{J}_i = e^{\frac{\pi \mathcal{J}_i}{2}}$, namely:

Lemma 1.7. (Gualtieri [19]) In a generalized Hermitian vector space one has

$$\star = -\mathbb{J}_1\mathbb{J}_2.$$

This means that we can read the decomposition of forms into SD and ASD from the $U^{p,q}$ decomposition, namely $\star|_{U^{p,q}} = i^{p+q}$. If we plot the (nontrivial) spaces $U^{p,q}$ in a lattice, each diagonal is made either of SD- or ASD-forms, with $U^{n,0}$ made of SD-forms (see Figure 1).

Definition 1.8. A *positive* $U(n)$ *structure* or a *positive Hermitian structure* on V is a generalized metric $\mathcal{G}$ and a complex structure $\mathcal{I}_+$ on V_+, the $+1$-eigenspace of $\mathcal{G}$, orthogonal with respect to the natural pairing and a *negative* $U(n)$ *structure* or *negative Hermitian structure* on V is a generalized metric with an orthogonal complex structure $\mathcal{I}_-$ on its -1-eigenspace. We say that a generalized complex structure $\mathcal{J}$ *extends* a positive/negative $U(n)$ structure $(\mathcal{G}, \mathcal{I})$ if $\mathcal{I}$ is the restriction of $\mathcal{J}$ to the appropriate space and $(\mathcal{G}, \mathcal{J})$ is a generalized Hermitian structure.

$$\begin{tikzcd}
 & & \text{SD} \arrow[dr, squiggly] & & & \\
 & \text{ASD} \arrow[dr, squiggly] & & U^{0,2} & & \\
\text{SD} \arrow[dr, squiggly] & & U^{-1,1} & & U^{1,1} & \\
 & U^{-2,0} & & U^{0,0} & & U^{2,0} \\
 & & U^{-1,-1} & & U^{1,-1} & \\
 & & & U^{0,-2} & &
\end{tikzcd} \tag{1.4}$$

Figure 1: Representation of the spaces of SD and ASD in terms of the (p,q)-decomposition of forms on a 4-dimensional generalized Hermitian structure.

Given a generalized Hermitian structure, since $\mathcal{J}_1$ and $\mathcal{G}$ commute, $\mathcal{J}_1$ preserves the eigenspaces of $\mathcal{G}$ and hence, upon restriction to $V_\pm$, one obtains a positive and a negative Hermitian structure. Conversely, a positive (resp. negative) Hermitian structure can be extended to $\mathbb{V}$ by declaring that it vanishes on V_- (resp. V_+). Then a pair of positive and negative Hermitian structures, $\mathcal{I}_+$, $\mathcal{I}_-$ gives rise to a generalized Hermitian structure by declaring that $\mathcal{J}_1 = \mathcal{I}_+ + \mathcal{I}_-$.

Given a positive $U(n)$ structure on V, we can use the isomorphism $V_+ \cong V$ to transport the metric and the complex structure $\mathcal{I}_+$ from V_+ to V, making it into a Hermitian vector space (V, g, I). Further, we can use $\mathcal{I}_+$ to define a complex structure $\mathcal{I}_-$ on V_- using the isomorphisms $V_+ \cong V \cong V_-$ and this way we have an extension of the $U(n)$ structure to a generalized Hermitian structure: namely we declare that $\mathcal{J}_1$ is $\mathcal{I}_+$ on V_+ and $\mathcal{I}_-$ on V_-, hence, in the metric splitting of $\mathbb{V}$, $\mathcal{J}_1$ is the generalized complex structure associated to the complex structure[2] I and consequenty $\mathcal{J}_2$ is the generalized complex structure associated to the Hermitian form $\omega = g \circ I$:

$$\mathcal{J}_1 = \begin{pmatrix} I & 0 \\ 0 & -I^* \end{pmatrix} \qquad \mathcal{J}_2 = \begin{pmatrix} 0 & -\omega^{-1} \\ \omega & 0 \end{pmatrix}. \tag{1.5}$$

For this set of choices, there is a relation between the (p,q)-decomposition of forms determined by the generalized Hermitian structure and the usual (p,q)-decomposition of forms determined by the complex structure I on V.

---

[2] Note that usually, the generalized complex structure associated to a given a complex structure I on a vector space differs from $\mathcal{J}_1$ by a sign, or said another way, by overall conjugation. Hence, all the usual concepts get conjugated with this definition, for example, later we will have $\partial_{\mathcal{J}_1} = \overline{\partial}$ and the generalized canonical bundle is $\wedge^{0,n}T^*M$

Proposition 1.9. (Cavalcanti [6]) *For $\mathcal{J}_1$ and $\mathcal{J}_2$ as above, the transformation*

$$\psi : \mathbb{V} \longrightarrow \mathbb{V} \qquad \psi(X+\xi) = e_*^{i\omega} e_*^{\frac{i}{2}\omega^{-1}}(X+\xi) \tag{1.6}$$

preserves $\mathcal{J}_1$ and maps $V \subset \mathbb{V}$ into the $+i$-eigenspace of $\mathcal{J}_2$ and $V^* \subset \mathbb{V}$ into the $-i$-eigenspace of $\mathcal{J}_2$. Therefore Ψ, the corresponding action on spinors, preserves the eigenspaces of $\mathcal{J}_1$ and maps $\wedge^k V^*$ into $U^{n-k}_{\mathcal{J}_2}$ that is

$$\begin{gathered}\Psi : \wedge^\bullet V^*_{\mathbb{C}} \longrightarrow \wedge^\bullet V^*_{\mathbb{C}} \quad \Psi(\varphi) = e^{i\omega} e^{\frac{i}{2}\omega^{-1}}(\varphi)\\ \Psi(\wedge^{p,q}V^*) = U^{q-p,n-p-q}_{\mathcal{J}_1,\mathcal{J}_2}.\end{gathered} \tag{1.7}$$

Proof. Since this statement is slightly different from that found in [6], we give a short sketch of the proof. Firstly, a simple computation gives an explicit expression for ψ:

$$\psi(X+\xi) = X - \frac{i}{2}\omega^{-1}(\xi) + \frac{\xi}{2} - i\omega(X).$$

Then the first claim is proved by restricting ψ to V and to V^*. Indeed, if we let L_2 denote the $+i$-eigenspace of $\mathcal{J}_2$ then

$$\begin{aligned}\psi(X) &= X - i\omega(X) \in L_2 \qquad \forall X \in V_{\mathbb{C}};\\ \psi(\xi) &= \frac{1}{2}(\xi - i\omega^{-1}(\xi)) \in \overline{L_2} \qquad \forall \xi \in V^*_{\mathbb{C}}\end{aligned} \tag{1.8}$$

Since ω is of type $(1,1)$ it maps $V^{0,1}$ into $V^{*0,1}$ hence it preserves the $\pm i$-eigenspaces of $\mathcal{J}_1$.

Finally, since the decomposition of forms associated to the bialgebroid $V, V^* \subset \mathbb{V}$ is given by the usual degree of forms, we see that the map above maps degree k forms into the space U^{n-k} for the symplectic structure. □

A positive Hermitian structure is fully determined by $V^{1,0}_+$, the $+i$-eigenspace of $\mathcal{I}_+$, that is, a positive $U(n)$ structure is a subspace $V^{1,0}_+ \subset \mathbb{T}_{\mathbb{C}}M$ for which

- $\dim(V^{1,0}) = n$;
- $V^{1,0}_+$ is isotropic with respect to the natural pairing;
- $V^{1,0}_+ \cap \overline{V^{1,0}_+} = \{0\}$ and
- $\langle v, \overline{v}\rangle > 0$ for all $v \in V^{1,0}_+ \backslash \{0\}$.

If we are given a generalized complex extension $\mathcal{J}$ of a positive Hermitian structure $(\mathcal{G}, \mathcal{I}_+)$ we obtain a bigrading of forms into $U^{p,q}$ as explained earlier. However, from the point of view of the $U(n)$ structure, the natural spaces to consider are $W^k_+ = \sum_{p+q=k} U^{p,q}$. Indeed W^n_+ corresponds to the vector space of all forms that are annihilated by the Clifford action of $V^{1,0}_+$ and $W^{n-2k}_+ = \wedge^k V^{0,1}_+ \cdot W^n_+$, that is, the spaces W^k_+ are solely determined by the complex structure $\mathcal{I}_+$. It is worth calling attention to the fact that W^k_+ is only nontrivial if $-n \leq k \leq n$ and $n-k=0 \mod 2$. Further, since the spaces W_+^k are the diagonals of the $U^{p,q}$ decomposition, each W_+^k is made of either SD- or ASD-forms, with W_+^n being SD.

Another description of the spaces $W_+^\bullet$ is obtained by extending $\mathcal{I}_+$ to an endomorphism of $\mathbb{T}M$ obtained by declaring that $\mathcal{I}_+|_{V_-}$ vanishes, so that $\mathcal{I}_+$ is a skew-symmetric operator on $\mathbb{T}M$ with respect to the natural pairing, that is, $\mathcal{I}_+ \in \mathfrak{spin}(\mathbb{T}M)$. Similarly to a generalized complex structure, $\mathcal{J}$, for which the space U^k is the ik-eigenspace of the action of $\mathcal{J}$, letting $\mathcal{I}_+$ act on forms one sees that the space W_+^k is the $i\frac{k}{2}$-eigenspace of $\mathcal{I}_+$. Hence, Lemma 1.7 takes the following form for positive $U(n)$ structures:

Lemma 1.10. Let $(\mathcal{G}, \mathcal{I}_+)$ be a positive Hermitian structure on V, and let $\mathbb{I}_+ = e^{\frac{\pi \mathcal{I}_+}{2}}$. Then

$$\mathbb{I}_+^2 = -\star,$$

that is $\star|_{W_+^k} = -i^k$.

Similarly, for a negative Hermitian structure $(\mathcal{G}, \mathcal{I}_-)$, we can extend $\mathcal{I}_-$ to $\mathbb{V}$ by declaring that it vanishes on V_+. If $\mathcal{J}_1$ is a generalized complex extension of $\mathcal{I}_-$, then the eigenspaces of $\mathcal{I}_-$ correspond to the anti-diagonals of the $U^{p,q}$ decomposition: $\mathcal{W}_-^k = \sum_{p-q=k} U^{p,q}$.

Finally we observe that the spaces $W_\pm^k$ have both even and odd forms, so one can refine this grading to a $\mathbb{Z} \times \mathbb{Z}_2$-grading:

$$W_\pm^{k,0} = W_\pm^k \cap (\wedge^{ev} V^* \otimes \mathbb{C}), \qquad W_\pm^{k,1} = W_\pm^k \cap (\wedge^{od} V^* \otimes \mathbb{C}). \tag{1.9}$$

In what follows we will refer to both spaces $W_\pm^k$ and $W_\pm^{k,l}$, with the understanding that if the $\mathbb{Z}_2$-grading is not particularly important, we will simply omit it.

2 Intrinsic torsion of generalized Hermitian structures

Except for a generalized metric, each of the structures introduced in Section 1 has an appropriate integrability condition. We let (M^{2n}, H) be a manifold endowed with a real closed 3-form H and consider the Courant bracket on sections of $\mathbb{T}M = TM \oplus T^*M$:

$$[\![X+\xi, Y+\eta]\!]_H = [X,Y] + \mathcal{L}_X \eta - i_Y d\xi - i_Y i_X H.$$

We will omit the 3-form from the bracket if it is clear from the context.

The Courant bracket is the derived bracket associated to the operator d^H, i.e., the following identity holds for all $v_1, v_2 \in \Gamma(\mathbb{T}M)$ and $\varphi \in \Omega^\bullet(M)$:

$$\begin{aligned} [\![v_1, v_2]\!]_H \cdot \varphi &= \{\{v_1, d^H\}, v_2\}\varphi \\ &= d^H(v_1 \cdot v_2 \cdot \varphi) + v_1 \cdot d^H(v_2 \cdot \varphi) - v_2 \cdot d^H(v_1 \cdot \varphi) - v_2 \cdot v_1 \cdot d^H \varphi, \end{aligned} \tag{2.1}$$

where $\cdot$ denotes the Clifford action of $\mathrm{Clif}(\mathbb{T}M)$ on $\wedge^\bullet T^*M$ and $\{\cdot,\cdot\}$ denotes the graded commutator of operators. We will use general properties of the derived bracket throughout these notes, and refer the reader to [25] for a concise presentation of its main properties.

The orthogonal action of a 2-form $B \in \Omega^2(M)$ on $\mathbb{T}M$ relates different Courant brackets:

$$[\![e_*^B v_1, e_*^B v_2]\!]_H = e_*^B [\![v_1, v_2]\!]_{H+dB}.$$

Definition 2.1. For each of the structures introduced in Definitions 1.6 and 1.8, we refer to the smooth assignment of such structure to T_xM for each $x \in M$ by including the adjective *almost* in the name of the structure.

Definition 2.2 (Integrability conditions)**.**

- A *generalized complex structure* is a generalized almost complex structure $\mathcal{J}$ whose $+i$-eigenspace is involutive with respect to the Courant bracket.
- A *generalized Hermitian structure* is a pair $(\mathcal{G}, \mathcal{J}_1)$ of generalized metric and compatible integrable generalized complex structure.
- A *positive (resp. negative) SKT structure* is a positive (resp. negative) almost $U(n)$ structure $(\mathcal{G}, \mathcal{I})$ for which the $+i$-eigenspace of $\mathcal{I}$ is involutive.
- A generalized (almost) complex structure $\mathcal{J}$ *extends* an SKT structure if $\mathcal{J}$ is fiberwise an extension of $\mathcal{I}$, in which case we say that $\mathcal{J}$ is a *generalized (almost) complex/Hermitian extension* of the SKT structure.
- A *generalized Kähler structure* is a generalized Hermitian structure $(\mathcal{G}, \mathcal{J}_1)$ for which $\mathcal{J}_1$ and $\mathcal{J}_2 = \mathcal{J}_1\mathcal{G}$ are integrable.

2.1 The Nijenhuis tensor

Let us spend some time to understand the *Nijenhuis tensor* of an almost generalized complex structure $\mathcal{J}$. This tensor is defined in the usual way, namely if L is the $+i$-eigenspace of $\mathcal{J}$

$$\mathrm{Nij} : \Gamma(\overline{L}) \times \Gamma(\overline{L}) \longrightarrow \Gamma(L); \qquad \mathrm{Nij}(X,Y) = -[\![X,Y]\!]^L, \tag{2.2}$$

where $\cdot^L$ indicates projection onto L. We can alternatively use the identification $L = \overline{L}^*$ from (1.3) to consider the operator

$$\begin{aligned} N : \Gamma(\overline{L}) \times \Gamma(\overline{L}) \times \Gamma(\overline{L}) &\longrightarrow \Omega^0(M;\mathbb{C}); \\ N(X,Y,Z) = -2\langle [\![X,Y]\!], Z\rangle &= 2\langle \mathrm{Nij}(X,Y), Z\rangle. \end{aligned} \tag{2.3}$$

As usual, Nij is a tensor, indeed, for $f \in \mathbb{C}^\infty(M;\mathbb{C})$ we have

$$\mathrm{Nij}(X, fY) = -[\![X, fY]\!]^L = -(f[\![X,Y]\!] + (\mathcal{L}_{\pi_T(X)} f)Y)^L = -f[\![X,Y]\!]^L = f\mathrm{Nij}(X,Y).$$

Further, the tensor $N \in \wedge^2 L \otimes L$ defined above actually lies in $\wedge^3 L$. Indeed, for $X, Y, Z \in \Gamma(\overline{L})$ we have

$$0 = \mathcal{L}_{\pi_T(X)}\langle Y, Z\rangle = -(\langle [\![X,Y]\!], Z\rangle + \langle Y, [\![X,Z]\!]\rangle) = \tfrac{1}{2}(N(X,Y,Z) + N(X,Z,Y)),$$

which shows that N is fully skew.

A different way to understand N arises by using the U^k decomposition determined by $\mathcal{J}$. Namely, letting $\mathcal{U}^k = \Gamma(U^k)$ (throughout the paper we denote the sheaf of sections of U^k by $\mathcal{U}^k$ and the sheaf of sections of $U^{p,q}$ by $\mathcal{U}^{p,q}$), one has:

Lemma 2.3. In an generalized almost complex manifold

$$d^H : \mathcal{U}^k \longrightarrow \mathcal{U}^{k-3} \oplus \mathcal{U}^{k-1} \oplus \mathcal{U}^{k+1} \oplus \mathcal{U}^{k+3}. \tag{2.4}$$

Further the map $\pi_{k+3} \circ d^H : \mathcal{U}^k \longrightarrow \mathcal{U}^{k+3}$ corresponds to the Clifford action of N, the Nijenhuis tensor defined in (2.3).

Proof. We prove first that

$$d^H : \mathcal{U}^k \longrightarrow \sum_{j \geq k-3} \mathcal{U}^j \tag{2.5}$$

and will do so by induction, starting at $k = n+1$ and working our way down. For $k = n+1$ we have that $U^{n+1} = \{0\}$ and the claim follows trivially.

Next we assume the result to be true for all $j > k$ and let $\rho \in \mathcal{U}^k$. For $v_1, v_2 \in \Gamma(L)$, using (2.1) we have

$$v_2 \cdot v_1 \cdot d^H \rho = -[\![v_1, v_2]\!] \cdot \rho + d^H(v_1 \cdot v_2 \cdot \rho) + v_1 d^H(v_2 \cdot \rho) - v_2 d^H(v_1 \cdot \rho).$$

Since the Clifford action of v_i sends $\mathcal{U}^j$ to $\mathcal{U}^{j+1}$, the inductive hypothesis implies that the last three terms lie in $\oplus_{j \geq k-1} \mathcal{U}^j$, while the first term, being the action of an element of $L \oplus \overline{L}$ on ρ lies in $\mathcal{U}^{k+1} \oplus \mathcal{U}^{k-1}$. Therefore we conclude that

$$v_2 \cdot v_1 \cdot d^H \rho \in \oplus_{j \geq k-1} \mathcal{U}^j \qquad \text{for all } v_1, v_2 \in \Gamma(L)$$

and (2.5) follows.

As d^H is a real operator and $U^{-k} = \overline{U^k}$ conjugating (2.5) we have that

$$d : \mathcal{U}^k \longrightarrow \sum_{j \leq k-3} \mathcal{U}^j.$$

Furthermore, if U^k is made of even forms then U^{k+1} is made of odd forms and vice versa, we have that

$$d^H : \mathcal{U}^k \longrightarrow \sum_{\substack{-3 \leq j-k \leq 3 \\ j-k = 1 \bmod 2}} \mathcal{U}^j,$$

therefore proving (2.4).

Now we prove that $\pi_{k+3} \circ d^H$ corresponds to the Clifford action of N. Once again we use induction, this time starting at $k = -n-1$ and moving upwards. Since $\mathcal{U}^{-n-1} = \{0\}$, the claim is trivial there. Assume now that for $j < k$ we have proved that $\pi_{j+3} \circ d^H$ is the Clifford action of N. Let $\rho \in \mathcal{U}^k$ and $v_1, v_2 \in \Gamma(\overline{L})$. Then using (2.1) we have

$$\begin{aligned}
v_2 v_1 \pi_{k+3} \circ d^H \rho &= \pi_{k+1}(v_2 v_1 d^H \rho) \\
&= \pi_{k+1}(-[\![v_1, v_2]\!] \rho + d^H v_1 v_2 \rho + v_1 d^H v_2 \rho - v_2 d^H v_1 \rho) \\
&= N(v_1, v_2)\rho + \pi_{k+1} \circ d^H (v_1 v_2 \rho) + v_1 \pi_{k+2} \circ d^H v_2 \rho - v_2 \pi_{k+2} \circ d^H v_1 \rho \\
&= \{v_2, \{v_1, N\}\}\rho + N v_1 v_2 \rho + v_1 N v_2 \rho - v_2 N v_1 \rho \\
&= v_2 v_1 N \rho,
\end{aligned}$$

where in third equality we have used that the component of $[\![v_1, v_2]\!]\rho$ in $\mathcal{U}^{k+1}$ is given by the Clifford action of the L component of $[\![v_1, v_2]\!]$, that is $\iota_{v_2}\iota_{v_1}N$ and in the fourth equality we used the inductive hypothesis as well as the fact that when acting on forms, the interior product of $v \in \Gamma(\overline{L})$ with $\varphi \in \Gamma(\wedge^\bullet L)$ is given by $\varphi(v)\rho = \{v, \varphi\}\rho$. The last equality follows by expanding the graded commutator and canceling out similar terms. □

If we compose $d^H|_{\mathcal{U}^k}$ with projection onto $\mathcal{U}^{k+1}$ and $\mathcal{U}^{k-1}$ we get operators

$$\partial_{\mathcal{J}} = \pi_{k+1} \circ d^H : \mathcal{U}^k \longrightarrow \mathcal{U}^{k+1} \qquad \text{and} \qquad \overline{\partial}_{\mathcal{J}} = \pi_{k-1} \circ d^H : \mathcal{U}^k \longrightarrow \mathcal{U}^{k-1},$$

and if the generalized complex structure $\mathcal{J}$ is clear from the context, we denote these operators simply by ∂ and $\overline{\partial}$. As proved by Gualtieri in [21], integrability is equivalent to the requirement

$$d^H : \mathcal{U}^k \longrightarrow \mathcal{U}^{k-1} \oplus \mathcal{U}^{k+1}; \qquad d^H = \partial + \overline{\partial},$$

so we can also see from this point of view how the vanishing of the Nijenhuis tensor implies integrability.

Since $(d^H)^2 = 0$, the operators ∂, $\overline{\partial}$, N, and $\overline{N}$ satisfy some relations.

Corollary 2.4. In an generalized almost complex manifold, the following relations and their complex conjugate hold

$$N^2 = 0, \qquad \{N, \partial\} = 0, \qquad \{N, \overline{\partial}\} + \partial^2 = 0, \qquad \{\partial, \overline{\partial}\} + \{N, \overline{N}\} = 0,$$

where $\{a, b\} = ab + (-1)^{|a||b|+1}ba$ is the graded commutator of the operators a and b, which, since all the operators above are odd, becomes simply $ab + ba$.

2.2 The intrinsic torsion and the road to integrability

With this understanding of the Nijenhuis tensor, we can give a pictorial description of the long road to integrability from almost generalized Hermitian to generalized Kähler. Indeed, given an almost generalized Hermitian structure, we get a splitting of forms into spaces $U^{p,q}$. According to Lemma 2.3, d^H can not change either the 'p' or the 'q' grading by more than three and it must switch parity. Hence d^H decomposes as a sum of eight operators and their complex conjugates

$$d^H = \delta_+ + \overline{\delta}_+ + \delta_- + \overline{\delta}_- + N_+ + \overline{N_+} + N_- + \overline{N_-} + N_1 + \overline{N_1} + N_2 + \overline{N_2} + N_3 + \overline{N_3} + N_4 + \overline{N_4};$$

$$\begin{aligned} \delta_+ &: \mathcal{U}^{p,q} \longrightarrow \mathcal{U}^{p+1,q+1}, & \delta_- &: \mathcal{U}^{p,q} \longrightarrow \mathcal{U}^{p+1,q-1}, \\ N_+ &: \mathcal{U}^{p,q} \longrightarrow \mathcal{U}^{p+3,q+3}, & N_- &: \mathcal{U}^{p,q} \longrightarrow \mathcal{U}^{p+3,q-3}, \\ N_1 &: \mathcal{U}^{p,q} \longrightarrow \mathcal{U}^{p-1,q+3}, & N_2 &: \mathcal{U}^{p,q} \longrightarrow \mathcal{U}^{p+1,q+3}, \\ N_3 &: \mathcal{U}^{p,q} \longrightarrow \mathcal{U}^{p+3,q+1}, & N_4 &: \mathcal{U}^{p,q} \longrightarrow \mathcal{U}^{p+3,q-1}. \end{aligned} \tag{2.6}$$

and we can draw in a diagram all the possible nontrivial components of $d^H|_{\mathcal{U}^{p,q}}$ as arrows (see Figure 2).

Once we require that $\mathcal{J}_1$ is integrable, i.e. we are dealing in fact with a generalized Hermitian structure, then d^H only changes the 'p' degree by ± 1 and several components

$$\begin{tikzcd}
\mathcal{U}^{p-3,q+3} & \mathcal{U}^{p-1,q+3} & & \mathcal{U}^{p+1,q+3} & \mathcal{U}^{p+3,q+3} \\
\mathcal{U}^{p-3,q+1} & \mathcal{U}^{p-1,q+1} & & \mathcal{U}^{p+1,q+1} & \mathcal{U}^{p+3,q+1} \\
& & \mathcal{U}^{p,q} \arrow[uull, bend left, "\overline{N_-}"'] \arrow[uul, bend right, "N_1"'] \arrow[uur, bend left, "N_2"] \arrow[uurr, bend right, "N_+"] \arrow[ull, "\overline{N_4}"] \arrow[ul, "\overline{\delta}_-"] \arrow[ur, "\delta_+"'] \arrow[urr, "N_3"'] \arrow[dll, "\overline{N_3}"'] \arrow[dl, "\overline{\delta}_+"'] \arrow[dr, "\delta_-"] \arrow[drr, "N_4"] \arrow[ddll, bend right, "\overline{N_+}"] \arrow[ddl, bend left, "\overline{N_2}"] \arrow[ddr, bend right, "\overline{N_1}"'] \arrow[ddrr, bend left, "N_-"'] & & \\
\mathcal{U}^{p-3,q-1} & \mathcal{U}^{p-1,q-1} & & \mathcal{U}^{p+1,q-1} & \mathcal{U}^{p+3,q-1} \\
\mathcal{U}^{p-3,q-3} & \mathcal{U}^{p-1,q-3} & & \mathcal{U}^{p+1,q-3} & \mathcal{U}^{p+3,q-3}
\end{tikzcd}$$

Figure 2: Representation of the nontrivial components of d^H when restricted to $\mathcal{U}^{p,q}$ for an generalized almost Hermitian structure.

$$\begin{tikzcd}
\mathcal{U}^{p-1,q+3} & & \mathcal{U}^{p+1,q+3} \\
\mathcal{U}^{p-1,q+1} & & \mathcal{U}^{p+1,q+1} \\
& \mathcal{U}^{p,q} \arrow[uul, "N_1"'] \arrow[uur, "N_2"] \arrow[ul, "\overline{\delta}_-"'] \arrow[ur, "\delta_+"] \arrow[dl, "\overline{\delta}_+"] \arrow[dr, "\delta_-"'] \arrow[ddl, "\overline{N_2}"'] \arrow[ddr, "\overline{N_1}"] & \\
\mathcal{U}^{p-1,q-1} & & \mathcal{U}^{p+1,q-1} \\
\mathcal{U}^{p-1,q-3} & & \mathcal{U}^{p+1,q-3}
\end{tikzcd}$$

Figure 3: Representation of the nontrivial components of d^H for a generalized Hermitian structure.

of d^H present in the nonintegrable case, now vanish and the diagram from Figure 2 clears up to the one presented in Figure 3.

Finally, if we require that $\mathcal{J}_2$ is also integrable, and hence we are in fact dealing with a generalized Kähler structure, the last two components of the Nijenhuis tensor, labeled N_1 and N_2 above, vanish and d^H decomposes as a sum of four operators, as pictured in Figure 4.

This shows that the obstruction for a generalized almost Hermitian structure to be a generalized Kähler structure is given by the tensors N_i, $i = 1, 2, 3, 4$ and $N_\pm$, that is, these tensors are the different components of the intrinsic torsion of a generalized almost Hermitian structure.

$$\begin{tikzcd}
\mathcal{U}^{p-1,q+1} & & \mathcal{U}^{p+1,q+1} \\
& \mathcal{U}^{p,q} \arrow[ul, "\overline{\delta}_-"'] \arrow[ur, "\delta_+"] \arrow[dl, "\overline{\delta}_+"] \arrow[dr, "\delta_-"'] & \\
\mathcal{U}^{p-1,q-1} & & \mathcal{U}^{p+1,q-1}
\end{tikzcd}$$

Figure 4: Representation of the nontrivial components of d^H for a generalized Kähler structure.

Definition 2.5. The *intrinsic torsion* of a generalized Hermitian manifold are the tensors N_i, $i = 1, 2, 3, 4$ and $N_\pm$.

3 SKT structures

In this section we provide properties equivalent to integrablity of an SKT structure as defined in 2.2. We will see that an SKT structure lies precisely half way between a generalized Hermitian and a generalized Kähler structure.

Firstly, we observe that the SKT condition can also be phrased in terms of the vanishing of components of the intrinsic torsion. Indeed, in the presence of an extension of, say, a positive $U(n)$ structure $(\mathcal{G}, \mathcal{I}_+)$ to a generalized almost complex structure $\mathcal{J}_1$ if we let $V_+^{1,0}$ be the $+i$-eigenspace of $\mathcal{I}_+$ then involutivity is equivalent to

$$\langle [\![v_1, v_2]\!], w_1 \rangle = 0 \qquad \text{for all } v_1, v_2 \in \Gamma(V_+^{1,0}), w \in \Gamma(V_+^{1,0} \oplus V_-),$$

that is, the components N_2, N_3 and N_+ of the intrinsic torsion vanish. A similar argument gives a characterization of negative SKT structures.

Proposition 3.1. *Let $\mathcal{G}$ be a generalized metric on a manifold with 3-form (M, H) and let $\mathcal{I}_\pm$ be a pair of positive and negative Hermitian structures compatible with $\mathcal{G}$. If $\mathcal{J}_1 = \mathcal{I}_+ + \mathcal{I}_-$ then $(\mathcal{G}, \mathcal{I}_+)$ is an SKT structure if and only if N_2, N_3 and N_+ vanish and $(\mathcal{G}, \mathcal{I}_-)$ is a negative SKT structure if and only if N_1, N_4 and N_- vanish.*

Corollary 3.2 (Gualtieri [20]). Let $\mathcal{G}$ be a generalized metric on a manifold with 3-form (M, H), $\mathcal{I}_\pm$ be a pair of positive and negative SKT structures compatible with $\mathcal{G}$ and $\mathcal{J}_1 = \mathcal{I}_+ + \mathcal{I}_-$. Then $(\mathcal{G}, \mathcal{J}_1)$ is a generalized Kähler structure.

Proof. According to Proposition 3.1, under the hypothesis, all components of the intrinsic torsion vanish hence $(\mathcal{G}, \mathcal{J}_1)$ is a generalized Kähler structure. □

Next, we describe the integrability condition for an SKT structure in terms of the decomposition of forms into $W_\pm^k$ and $U^{p,q}$ (for a fixed generalized complex extension $\mathcal{J}_1$) described in the previous section. Throughout the paper we denote the sheaf of sections of $W_\pm^k$ by $\mathcal{W}_\pm^k$ and the sheaf of sections of $W_\pm^{k,l}$ by $\mathcal{W}_\pm^{k,l}$.



Figure 5: Representation of the nontrivial components of d^H for a positive SKT structure



Figure 6: Representation of the nontrivial components of d^H for a negative SKT structure

Theorem 3.3. *Let $(\mathcal{G}, \mathcal{I}_+)$ be a positive almost Hermitian structure on a manifold with 3-form (M^{2n}, H). The the following are equivalent:*

1. *$(\mathcal{G}, \mathcal{I}_+)$ is a positive SKT structure;*
2. *$d^H : \mathcal{W}^k_+ \longrightarrow \mathcal{W}^{k-2}_+ \oplus \mathcal{W}^k_+ \oplus \mathcal{W}^{k+2}_+$ for all k;*
3. *$d^H : \mathcal{W}^n_+ \longrightarrow \mathcal{W}^{n-2}_+ \oplus \mathcal{W}^n_+$.*

Proof. One can give a direct proof of this result using only the positive almost Hermitian data with an argument completely analogous to that already given in Lemma 2.3. Yet, since we have already developed the tools for generalized almost Hermitian structures, we follow a less intrinsic route.

We will first prove that 1) implies 2). Let $\mathcal{I}_-$ be any complex structure on V_- orthogonal with respect to the natural pairing so that $\mathcal{G}$ and $\mathcal{J}_1 = \mathcal{I}_+ + \mathcal{I}_-$ form a generalized almost Hermitian structure. Then according to Proposition 3.1, N_2, N_3 and N_+ vanish. Then, we see that d^H splits into three components:

$$d^H = \delta_+^N + \overline{\delta_+^N} + \not\delta_- \tag{3.1}$$

$$\begin{array}{lll}
\not\delta_- \colon \mathcal{W}_+^k \longrightarrow \mathcal{W}_+^k; & \qquad & \not\delta_- = \delta_- + \overline{\delta}_- + N_- + \overline{N_-}, \\
\delta_+^N \colon \mathcal{W}_+^k \longrightarrow \mathcal{W}_+^{k+2}; & \qquad & \delta_+^N = \delta_+ + N_1 + N_4, \\
\overline{\delta_+^N} \colon \mathcal{W}_+^k \longrightarrow \mathcal{W}_+^{k-2}; & \qquad & \overline{\delta_+^N} = \overline{\delta}_+ + \overline{N_1} + \overline{N_4}.
\end{array}$$

Condition 2) clearly implies condition 3). Finally to prove that 3) implies 1) we once again choose a complex structure $\mathcal{I}_-$ on V_- and observe that since $\overline{N_2}, \overline{N_3}$ and $\overline{N_+}$ are tensors, it is enough to check that they vanish when applied to spaces where their action is effective. But for $\varphi \in \mathcal{U}^{n,0} \subset \mathcal{W}_+^n$, the components of d^H landing on $\mathcal{W}_+^{n-6}$ and $\mathcal{W}_+^{n-4}$ are precisely $\overline{N_+} \cdot \varphi$ and $\overline{N_3} \cdot \varphi$ and the vanishing of these forms for $\varphi \neq 0$ implies that $\overline{N_+} = \overline{N_3} = 0$. Similarly, if $\psi \in \mathcal{U}^{0,n}$, the $\mathcal{W}_+^{n-4}$ component of d^H is $\overline{N_2} \cdot \psi$, hence the vanishing of this component implies that $\overline{N_2} = 0$ and Proposition 3.1 implies that $\mathcal{I}_+$ is integrable. □

While the different $U^{p,q}$ components of d^H obtained in terms of the generalized almost Hermitian extension of the SKT structure depend on the particular extension chosen, the operators δ_+^N, $\overline{\delta_+^N}$ and $\not\delta_-$ depend only on the SKT data, since they are the decomposition of d^H obtained from the eigenspaces of $\mathcal{I}_+$.

Corollary 3.4. The following hold

$$(\delta_+^N)^2 = (\overline{\delta_+^N})^2 = 0; \qquad \{\delta_+^N, \not\delta_-\} = 0; \qquad \{\overline{\delta_+^N}, \not\delta_-\} = 0; \qquad \{\delta_+^N, \overline{\delta_+^N}\} + (\not\delta_-^H)^2 = 0;$$

where $\{\cdot, \cdot\}$ indicates the graded commutator of the operators.

In the arguments up to now, given a positive SKT structure, we have chosen the complex structure $\mathcal{I}_-$ rather freely, but one can still require some integrability of $\mathcal{I}_-$ without restricting the geometry, as we argue next.

Proposition 3.5. (Cavalcanti [7]) *Given a positive SKT structure* $(\mathcal{G}, \mathcal{I}_+)$ *on* (M, H), *there is a generalized complex structure* $\mathcal{J}_1$ *of complex type which extends* $\mathcal{I}_+$. *Further* $\mathcal{J}_2 = \mathcal{G}\mathcal{J}_1$ *is of symplectic type and its canonical bundle has nonvanishing section* ρ_2 *such that* $\overline{\partial}_{\mathcal{J}_1}\rho_2 = 0$.

Idea of the proof. Indeed, we can use the procedure from Section 1 to produce an almost complex structure I with compatible metric g on M and we can extend the SKT structure to an almost Hermitian structure $(\mathcal{G}, \mathcal{J}_1)$, where $\mathcal{J}_1$ is the generalized complex structure corresponding to the complex structure I, as in (1.5). We let $\omega = g \circ I$ be the corresponding Hermitian form. Then integrability of the SKT structure is equivalent to the integrability of I (and hence of $\mathcal{J}_1$) and the condition $d^c\omega = H$, where H is the 3-form associated to the metric splitting of $\mathbb{T}M$. In particular $\mathcal{J}_1$ is a generalized complex extension of the SKT structure. Finally, the canonical bundle of $\mathcal{J}_2$ is generated by $\rho_2 = e^{i\omega}$ and

$$d^H e^{i\omega} = e^{i\omega} \wedge (H + id\omega) = 2ie^{i\omega}\overline{\partial}\omega \in \mathcal{U}^1_{\mathcal{J}_1},$$

hence $\overline{\partial}_{\mathcal{J}_1}\rho_2 = 0$. ☐

Theorem 3.6. *Let (M, H) be a manifold with 3-form and $(\mathcal{G}, \mathcal{J}_1)$ be an generalized almost Hermitian structure on M. Then the following are equivalent:*

1. *$(\mathcal{G}, \mathcal{J}_1)$ is an integrable generalized complex extension of a positive SKT structure;*
2. *$d^H : \mathcal{U}^{p,q} \longrightarrow \mathcal{U}^{p+1,q+1} \oplus \mathcal{U}^{p+1,q-1} \oplus \mathcal{U}^{p+1,q-3} \oplus \mathcal{U}^{p-1,q+3} \oplus \mathcal{U}^{p-1,q+1} \oplus \mathcal{U}^{p-1,q-1}$ for all $p, q \in \mathbb{Z}$;*
3. *$d^H : \mathcal{U}^{p,n-p} \longrightarrow \mathcal{U}^{p+1,n-p-1} \oplus \mathcal{U}^{p+1,n-p-3} \oplus \mathcal{U}^{p-1,n-p+1} \oplus \mathcal{U}^{p-1,n-p-1}$ for all $p \in \mathbb{Z}$;*
4. *$d^H : \mathcal{U}^{0,n} \longrightarrow \mathcal{U}^{-1,n-1} \oplus \mathcal{U}^{1,n-1} \oplus \mathcal{U}^{1,n-3}$ and $d^H : \mathcal{U}^{n,0} \longrightarrow \mathcal{U}^{n-1,1} \oplus \mathcal{U}^{n-1,-1}$.*

Proof. The fact that 1) implies 2) follows from Proposition 3.1 and integrability of $\mathcal{J}_1$. The implications 2) ⇒ 3) ⇒ 4) are immediate. Finally, similarly to the proof of Theorem 3.3, the values of the different components of the intrinsic torsion are fully determined by their action on $\mathcal{U}^{n,0}$ and $\mathcal{U}^{0,n}$ and 4) implies the vanishing of all components of the intrinsic torsion except from N_1. ☐

In light of Proposition 3.5, this theorem allows us to define six differential operators on a generalized complex extension of a positive SKT structure:

$$\begin{gathered} d^H = \delta_- + \overline{\delta}_- + N + \overline{N} + \delta_+ + \overline{\delta}_+ \qquad (3.2) \\ \delta_- : \mathcal{U}^{p,q} \longrightarrow \mathcal{U}^{p+1,q-1}; \qquad \delta_+ : \mathcal{U}^{p,q} \longrightarrow \mathcal{U}^{p+1,q+1}; \qquad N : \mathcal{U}^{p,q} \longrightarrow \mathcal{U}^{p-1,q+3}. \end{gathered}$$

We see that $N = N_1$ is the Nijenhuis tensor of $\mathcal{J}_2$, in particular, N is tensorial and this theorem shows that if $(\mathcal{G}, \mathcal{J}_1)$ is a generalized Hermitian extension of an SKT structure then half of the Nijenhuis tensor of $\mathcal{J}_2$ vanishes, namely the component N_2 from (2.6). Also, this decomposition allows us to express the operators δ_+^N and $\not\delta_-$ from (3.1) in terms of δ_+, δ_-, N and their conjugates:

$$\delta_+^N = \delta_+ + N; \qquad \not\delta_- = \delta_- + \overline{\delta}_-. \qquad (3.3)$$

Since $(d^H)^2 = 0$ these operators satisfy a number of relations.

Corollary 3.7. The following relations and their complex conjugates hold:

$$\delta_+^2 = N^2 = 0; \qquad \{\delta_+, N\} = 0;$$

$$\delta_-^2 = -\{\overline{N}, \delta_+\}; \qquad \{\overline{N}, \delta_-\} = 0; \qquad \{\delta_-, \delta_+\} = 0; \qquad \{\delta_+, \overline{\delta}_-\} = -\{\delta_-, N\};$$

$$\{\delta_+, \overline{\delta}_+\} + \{\delta_-, \overline{\delta}_-\} + \{N, \overline{N}\} = 0.$$

$$\begin{tikzcd}
\mathcal{U}^{p-1,q+3} & & \\
& \mathcal{U}^{p,q+2} & \\
\mathcal{U}^{p-1,q+1} & & \mathcal{U}^{p+1,q+1} \\
& \mathcal{U}^{p,q} \arrow[uuul, "N"'] \arrow[ul, "\overline{\delta}_-"] \arrow[dl, "\overline{\delta}_+"'] \arrow[ur, "\delta_+"'] \arrow[dr, "\delta_-"] \arrow[dddr, "\overline{N}"] & \\
\mathcal{U}^{p-1,q-1} & & \mathcal{U}^{p+1,q-1} \\
& \mathcal{U}^{p,q-2} & \\
& & \mathcal{U}^{p+1,q-3}
\end{tikzcd}$$

Figure 7: Decomposition of d^H for a generalized complex extension of a positive SKT structure.

Corollary 3.8. Let $(\mathcal{G},\mathcal{J}_1)$ be a generalized Hermitian structure on a manifold with 3-form (M,H) and let $\mathcal{J}_2=\mathcal{G}\mathcal{J}_1$ be the associated generalized almost complex structure. If the canonical bundle of $\mathcal{J}_2$ admits a $\overline{\partial}_{\mathcal{J}_1}$-closed trivialization, then $(\mathcal{G},\mathcal{J})$ is an extension of an SKT structure.

Proof. Since $\mathcal{J}_1$ is integrable, $d^H:\mathcal{U}^{n,0}\longrightarrow\mathcal{U}^{n-1,1}\oplus\mathcal{U}^{n-1,-1}$. Due to Lemma 2.3, we have that

$$d^H:\mathcal{U}^{0,n}\longrightarrow\mathcal{U}^{-1,n-3}\oplus\mathcal{U}^{-1,n-1}\oplus\mathcal{U}^{1,n-1}\oplus\mathcal{U}^{1,n-3}.$$

But according to the hypothesis there is a trivialization ρ of $U^{0,n}$ such that $\overline{\partial}_{\mathcal{J}_1}\rho=0$, that is, for such ρ the components of $d^H\rho$ lying in $\mathcal{U}^{-1,n-3}\oplus\mathcal{U}^{-1,n-1}$ vanish. In particular, the component in $\mathcal{U}^{-1,n-3}$ is part of the Nijenhuis tensor of $\mathcal{J}_2$ and since it vanishes on ρ it is the zero tensor. So we see that, in fact

$$d^H:\mathcal{U}^{0,n}\longrightarrow\mathcal{U}^{-1,n-1}\oplus\mathcal{U}^{1,n-1}\oplus\mathcal{U}^{1,n-3}.$$

and the last condition of Theorem (3.6) holds. ☐

The same results hold for *negative SKT* structures. One should bear in mind, however that the relevant spaces for a negative structure are given by $W_-^k=\sum_{p-q=k}U^{p,q}$, that is, each W_-^k is an antidiagonal and integrability is equivalent to

$$d^H:\mathcal{W}_-^k\longrightarrow\mathcal{W}_-^{k+2}\oplus\mathcal{W}_-^k\oplus\mathcal{W}_-^{k-2},$$

or, in terms of the $U^{p,q}$ decomposition obtained by choosing a generalized complex extension, d^H decomposes as shown in Figure 8.

So an SKT structure (positive or negative) corresponds to a generalized Hermitian structure in which half of the Nijenhuis tensor of $\mathcal{J}_2$ vanishes.

$$\begin{tikzcd}
 & & \mathcal{U}^{p+1,q+3} \\
\mathcal{U}^{p-1,q+1} & & \mathcal{U}^{p+1,q+1} \\
 & \mathcal{U}^{p,q} \arrow[ruu, "N_2"] \arrow[lu, "\overline{\delta}_-"'] \arrow[ru, "\delta_+"'] \arrow[ld, "\overline{\delta}_+"] \arrow[rd, "\delta_-"] \arrow[ldd, "\overline{N_2}"'] & \\
\mathcal{U}^{p-1,q-1} & & \mathcal{U}^{p+1,q-1} \\
\mathcal{U}^{p-1,q-3} & &
\end{tikzcd}$$

Figure 8: Decomposition of d^H for a negative SKT structure.

4 Parallel Hermitian and bi-Hermitian structures

4.1 Parallelism and the intrinsic torsion

The existence of a relationship between connections with closed, skew symmetric torsion and the Courant bracket was made evident by Hitchin in [22]. Precisely, given a generalized metric on $\mathbb{T}M$, we let H be the 3-form corresponding to the metric splitting and g be the induced metric on M. Also, for $X \in \Gamma(TM)$ we let $X_\pm \in \Gamma(V_\pm)$ be unique lifts of X to $V_\pm$, we let $\pi_\pm : \mathbb{T}M \longrightarrow V_\pm$ be the orthogonal projections onto $V_\pm$ and $\pi_T : \mathbb{T}M \longrightarrow TM$ be the natural projection.

Proposition 4.1 (Hitchin [22])**.** *Let* $\widetilde{\nabla}^\pm$ *be the unique metric connection whose torsion is skew symmetric and equal to* $\mp H$*. Then*

$$\widetilde{\nabla}^+_X Y = \pi_T \circ \pi_+ [\![X_-, Y_+]\!],$$

$$\widetilde{\nabla}^-_X Y = \pi_T \circ \pi_- [\![X_+, Y_-]\!].$$

From this proposition, we see that the isomorphisms $\pi_T : V_\pm \longrightarrow TM$ relate the connections with torsion $\mp H$ to the operators

$$\nabla^+ \colon \Gamma(V_-) \times \Gamma(V_+) \longrightarrow \Gamma(V_+); \qquad \nabla^+_v w = \pi_+ [\![v, w]\!], \quad v \in \Gamma(V_+), w \in \Gamma(V_-), \tag{4.1}$$

$$\nabla^- \colon \Gamma(V_+) \times \Gamma(V_-) \longrightarrow \Gamma(V_-); \qquad \nabla^-_w v = \pi_- [\![w, v]\!], \quad v \in \Gamma(V_+), w \in \Gamma(V_-). \tag{4.2}$$

As we will work with the spaces $V_\pm$ directly, we will use $\nabla^\pm$ instead of the connections they induce on TM, with the understanding that these are equivalent operators, so, for example, if $\widetilde{\nabla}^+$ has holonomy in $U(n)$, then M has an almost Hermitian structure (g, I) which is parallel with respect to $\widetilde{\nabla}^+$ and via the isomorphism $\pi_T : V_+ \longrightarrow TM$, V_+ gets a positive almost Hermitian structure which is parallel for ∇^+ so that ∇^+ has holonomy in $U(n)$ and the converse statement also holds.

Definition 4.2. A *parallel positive (resp. negative) Hermitian structure* is a positive (resp. negative) almost Hermitian structure which is parallel with respect to ∇^+, (resp. ∇^-). A *parallel bi-Hermitian structure* or *parallel* $U(n) \times U(n)$*-structure* is a triple (g, I_+, I_-) such that (g, I_+) is a parallel positive Hermitian structure and (g, I_-) is a parallel negative Hermitian structure.

Proposition 4.3. *If the connection* ∇^+ *has holonomy in* $U(n)$*, the corresponding positive almost Hermitian structure satisfies*

$$[\![\Gamma(V_+^{1,0}), \Gamma(V_+^{1,0})]\!] \subset \Gamma(V_+ \otimes \mathbb{C}). \tag{4.3}$$

And conversely, if a positive Hermitian structure satisfies (4.3)*, then it is parallel with respect to* ∇^+*.*

Similarly, ∇^- *has holonomy in* $U(n)$ *if and only if*

$$[\![\Gamma(V_-^{1,0}), \Gamma(V_-^{1,0})]\!] \subset \Gamma(V_- \otimes \mathbb{C}). \tag{4.4}$$

Proof. Let $\mathcal{I}_+$ be a complex structure on V_+ orthogonal with respect to the natural pairing and let $V_+^{1,0}$ be its $+i$-eigenspace. Then, for $v_1, v_2 \in \Gamma(V_+^{1,0})$ and $w \in \Gamma(V_-)$, we have that $\langle v_1, w\rangle = 0$, as V_+ and V_- are orthogonal with respect to the natural pairing. Hence

$$\begin{aligned} 0 &= \mathcal{L}_{\pi_T v_1}\langle w, v_2\rangle \\ &= \langle [\![v_1, w]\!], v_2\rangle + \langle w, [\![v_1, v_2]\!]\rangle \\ &= -\langle [\![w, v_1]\!], v_2\rangle + \langle w, [\![v_1, v_2]\!]\rangle \\ &= -\langle \nabla^+_w v_1, v_2\rangle + \langle w, [\![v_1, v_2]\!]\rangle. \end{aligned} \tag{4.5}$$

where in the third equality we used again that V_+ and V_- are orthogonal with respect to the natural pairing, and hence $[\![v_1, w]\!] = -[\![w, v_1]\!]$ and in the fourth equality we used that $v_2 \in V_+$ and hence the V_- component of $[\![w, v_1]\!]$ is annihilated by the natural pairing.

If $\mathcal{I}_+$ is parallel with respect to ∇^+, then $\langle \nabla^+_w v_1, v_2\rangle$ vanishes for all $v_1, v_2 \in \Gamma(V_+^{1,0})$ and $w \in \Gamma(V_-)$ and hence, according to (4.5), so does $\langle w, [\![v_1, v_2]\!]\rangle$, showing that $[\![v_1, v_2]\!]$ must be orthogonal to V_- and hence, a section of V_+. Conversely, if , $[\![v_1, v_2]\!] \in \Gamma(V_+)$, then for all $w \in \Gamma(V_-)$, $\langle w, [\![v_1, v_2]\!]\rangle = 0$ hence (4.5) implies that $\langle \nabla^+_w v_1, v_2\rangle = 0$ showing that $\nabla^+_w v_1$ is orthogonal to $V_+^{1,0}$. Since $V_+^{1,0}$ is a maximal isotropic of $V_+ \otimes \mathbb{C}$, we conclude that $\nabla^+_w v_1 \in \Gamma(V_+^{1,0})$ for all $w \in V_-$ and $v_1 \in \Gamma(V_+^{1,0})$, hence $\mathcal{I}_+$ is parallel with respect to ∇^+. ☐

Remark. Here we obtain, in a new light, a well known contrast between connections with torsion and the Levi–Civita connection regarding integrability. Indeed, for the Levi–Civita connection, reduction of the holonomy group to $U(n)$ implies integrability of the complex structure, but that is known not to be the case for connections with torsion. From our point of view, this is the difference between the reduced holonomy condition

$$[\![\Gamma(V_+^{1,0}), \Gamma(V_+^{1,0})]\!] \subset \Gamma(V_+)$$

and the SKT condition

$$[\![\Gamma(V_+^{1,0}), \Gamma(V_+^{1,0})]\!] \subset \Gamma(V_+^{1,0}).$$

Theorem 4.4. *Let (M, H) be a manifold with 3-form, $\mathcal{G}$ be a generalized metric on M and $\mathcal{I}_+$, $\mathcal{I}_-$ be a positive and a negative almost Hermitian structure on M*

- *$\mathcal{I}_+$ is parallel with respect to ∇^+ if and only if $N_2 = N_3 = 0$, i.e.,*

$$d^H : \mathcal{W}_+^k \longrightarrow \mathcal{W}_+^{k-6} \oplus \mathcal{W}_+^{k-2} \oplus \mathcal{W}_+^k \oplus \mathcal{W}_+^{k+2} \oplus \mathcal{W}_+^{k+6};$$

- *$\mathcal{I}_-$ is parallel with respect to ∇^- if and only if $N_1 = N_4 = 0$, i.e.,*

$$d^H : \mathcal{W}_-^k \longrightarrow \mathcal{W}_-^{k-6} \oplus \mathcal{W}_-^{k-2} \oplus \mathcal{W}_-^k \oplus \mathcal{W}_-^{k+2} \oplus \mathcal{W}_-^{k+6};$$

- *$(\mathcal{G}, \mathcal{I}_+, \mathcal{I}_-)$ is a parallel bi-Hermitian structure if and only if $N_i = 0$ for $i = 1, 2, 3, 4$.*

Proof. Since V_- is the orthogonal complement of V_+ with respect to the natural pairing, it is clear that (4.3) is equivalent to the following two conditions

$$\langle [\![v_1, v_2]\!], w\rangle = 0 \text{ for all } v_1, v_2 \in \Gamma(V_+^{1,0}), w \in \Gamma(V_-^{0,1}).$$

$$\langle [\![v_1, v_2]\!], w\rangle = 0 \text{ for all } v_1, v_2 \in \Gamma(V_+^{1,0}), w \in \Gamma(V_-^{1,0});$$

but the first condition is equivalent to the vanishing of $\overline{N_2}$ and the second, to the vanishing of $\overline{N_3}$. Finally, since the component of d^H mapping $\mathcal{W}^k$ to $\mathcal{W}^{k+4}$ is given by the sum $N_2 + N_3$, we see that the vanishing of this component is also equivalent to the parallel condition.

The remaining claims are proved similarly. □

The decomposition of d^H for an almost generalized Hermitian structure extending a parallel positive Hermitian structure is depicted in Figure 9 and the decomposition of d^H for a parallel bi-Hermitian structure is depicted in Figure 10.

The same argument used in Theorem 4.4, shows that the obstructions for parallel structures to be integrable are given by $N_\pm$:

Proposition 4.5. *Let (M, H) be a manifold with 3-form.*

- *A parallel positive (resp. negative) almost Hermitian structure is a positive (resp. negative) SKT structure if and only if $N_+ = 0$ (resp. $N_- = 0$);*
- *A parallel bi-Hermitian structure is a generalized Kähler structure if and only if $N_+ = N_- = 0$.*

Corollary 4.6. Let M^4 be a four dimensional manifold.

- A parallel positive/negative Hermitian structure is a positive/negative SKT structure;
- A parallel bi-Hermitian structure is a generalized Kähler structure.

Proof. Since M is four dimensional, $V_\pm^{1,0}$ are two dimensional complex vector spaces, so $\wedge^3 V_\pm^{1,0} = \{0\}$ and hence $N_\pm \in \wedge^3 V_\pm^{1,0}$ are the trivial tensors. □

We finish this section with a characterization of parallel structures similar to the differential geometric characterization of SKT structures.



Figure 9: Representation of the nontrivial components of d^H for an almost generalized Hermitian structure extending a parallel positive Hermitian structure



Figure 10: Representation of the nontrivial components of d^H for a parallel bi-Hermitian structure.

Theorem 4.7. *Let M be a manifold with 3-form, $\mathcal{G}$ be a generalized metric and $\mathcal{I}_+$, $\mathcal{I}_-$ be a positive and a negative almost Hermitian structure on M and $\mathcal{J}_1 = \mathcal{I}_+ + \mathcal{I}_-$. Let H be the 3-form corresponding to the metric splitting of $\mathbb{T}M$ and $(g, I_\pm)$ be the corresponding bi-Hermitian data. Then*

1. *$\mathcal{I}_+$ is a parallel positive Hermitian structure if and only if*

$$A_+ \in \otimes^3 T_+^{*1,0}M; \quad A_+(X,Y,Z) = \omega_+([X,Y],Z), \quad X,Y,Z \in \Gamma(T_+^{1,0}M)$$

is a skew symmetric tensor[3] *and*

$$d\omega_+ = I_+^* H;$$

where $T_+^{1,0}M$ *is the* $+i$*-eigenspace of* I_+, $\omega_+(X,Y) = g(I_+X,Y)$ *and* $I_+^*|_{\Omega^{p,q}} = i(p-q)$*;*

2. $(\mathcal{G}, \mathcal{J}_1)$ *is a parallel negative Hermitian structure if and only if*

$$A_- \in \otimes^3 T_-^{*1,0}M; \quad A_-(X,Y,Z) = \omega_-([X,Y],Z)$$

is a skew symmetric tensor and

$$d\omega_- = -I_-^* H;$$

where $T_-^{1,0}M$ *is the* $+i$*-eigenspace of* I_- *and* $\omega_-(X,Y) = g(I_-X,Y)$*;*

3. $(\mathcal{G}, \mathcal{J}_1)$ *is a parallel bi-Hermitian structure if and only if 1) and 2) hold.*

Proof. We will only prove 1) as 2) is completely analogous and 3) follows from 1) and 2).

Let $v, w \in \Gamma(V_+^{1,0})$ and let $X = \pi_T(v)$ and $Y = \pi_T(w) \in \Gamma(T_+^{1,0}M)$ so that $v = X + g(X)$ and $w = Y + g(Y)$, where $T_+^{1,0}M$ is the $+i$-eigenbundle of the almost complex structure I_+. Since $\omega_+ = g \circ I_+$, we have that $g(X) = -i\omega_+(X)$ and hence

$$\begin{aligned}
[\![v,w]\!]_H &= [\![X - i\omega_+(X), Y - i\omega_+(Y)]\!]_H \\
&= [\![e^{i\omega_+}X, e^{i\omega_+}Y]\!]_H \\
&= e^{i\omega_+}[\![X,Y]\!]_{H+id\omega_+} \\
&= [X,Y] - \iota_Y\iota_X(H + id\omega_+) - i\iota_{[X,Y]}\omega_+.
\end{aligned}$$

Therefore, $[\![v,w]\!]$ is a section of V_+ if and only if

$$g([X,Y]) = -\iota_Y\iota_X(H + id\omega_+) - i\iota_{[X,Y]}\omega_+. \tag{4.6}$$

Splitting $[X,Y]$ into its $(1,0)$ and $(0,1)$-components, we have $[X,Y] = [X,Y]^{1,0} - \mathrm{Nij}(X,Y) = [X,Y]^{1,0} - \{\iota_Y, \{\iota_X, N\}\}$, where Nij and $N \in \Gamma(\wedge^3 L_{I_+})$ are the tensors introduced in (2.2) and (2.3) for the generalized complex structure associated to I_+

$$\mathcal{J}_{I_+} = \begin{pmatrix} -I_+ & 0 \\ 0 & I_+^* \end{pmatrix}$$

computed with respect to the Courant bracket with zero 3-form. Similarly, $d\omega_+$ has four components:

$$d\omega_+ = N \cdot \omega_+ + \partial\omega_+ + \overline{\partial}\omega_+ + \overline{N} \cdot \omega_+.$$

Hence, (4.6) is equivalent to

$$\begin{aligned}
-i\iota_{[X,Y]^{1,0}}\omega_+ - i\{\iota_Y, \{\iota_X, N\}\}\omega_+ = &- \iota_Y\iota_X(H + i(N \cdot \omega_+ + \partial\omega_+ + \overline{\partial}\omega_+ + \overline{N} \cdot \omega_+)) + \\
&- i(\iota_{[X,Y]^{1,0}}\omega_+ - \{\iota_Y, \{\iota_X, N\}\}\omega_+).
\end{aligned}$$

---

[3] This condition is sometimes phrased in the literature as "the Nijenhuis tensor of I_+ is fully skew symmetric". Since we have a slightly different definition of Nijnehuis tensor and ours is always fully skew symmetric, we avoid the use of this common terminology to refer to this condition.

Since $\overline{\partial}\omega_+ \in \Omega^{1,2}(M)$ and $\overline{N}\cdot\omega_+ \in \Omega^{0,3}(M)$, these terms are annihilated by the interior product by $X, Y \in \Gamma(T_+^{1,0}M)$ and we obtain that (4.6) is equivalent to

$$i\iota_Y\iota_X H = \iota_Y\iota_X\partial\omega - 2\{\iota_Y, \{\iota_X, N\}\}\cdot\omega_+ + \iota_Y\iota_X N\cdot\omega_+. \tag{4.7}$$

Equating the $(0,1)$ components of both sides of (4.7), we get

$$i\iota_Y\iota_X H^{2,1} = \iota_Y\iota_X\partial\omega_+,$$

hence

$$iH^{2,1} = \partial\omega_+. \tag{4.8}$$

To determine the $(1,0)$ components of both sides of (4.7) we contract the whole equation with $Z \in T_+^{1,0}M$:

$$iH(X,Y,Z) = 2\{\iota_Y, \{\iota_X, N\}\}\cdot\iota_Z\omega_+ + \iota_Z\iota_Y\iota_X N\cdot\omega_+. \tag{4.9}$$

Since $H(X,Y,Z)$ and $N\cdot\omega_+$ are skew forms, this equality can only hold if

$$(X,Y,Z)\mapsto\{\iota_Y,\{\iota_X,N\}\}\cdot\iota_Z\omega_+$$

is also skew and, spelling it, out we see that the tensor above is nothing but A_+ from the statement of the theorem. Further, expanding the last term and using skew symmetry of A_+ we get

$$\iota_Z\iota_Y\iota_X N\cdot\omega_+ = -\{\iota_Y,\{\iota_X,N\}\}\iota_Z\omega + c.p = -3\{\iota_Y,\{\iota_X,N\}\}\iota_Z\omega.$$

Hence, (4.9) becomes

$$3iH(X,Y,Z) = \iota_Z\iota_Y\iota_X N\cdot\omega,$$

which is equivalent to

$$3iH^{3,0} = N\cdot\omega_+ \tag{4.10}$$

and (4.7) is equivalent to (4.8) and (4.10) an skew symmetry of A_+. Since H is real, we obtain that the parallel condition is equivalent to

$$I_+^*H = 3iH^{3,0} + iH^{2,1} - i\overline{H^{2,1}} - 3i\overline{H^{3,0}} = N\cdot\omega_+ + \partial\omega_+ + \overline{\partial}\omega_+ + \overline{N}\cdot\omega_+ = d\omega_+.$$

□

5 Hodge theory

In this section we develop Hodge theory for manifolds with parallel Hermitian, bi-Hermitian and SKT structures . Our main result is that for a parallel positive Hermitian structure the Laplacian preserves the spaces $\mathcal{W}_+^{k,l}$ and hence induces a decomposition of the d^H-cohomology accordingly. This is in contrast with the fact that for usual manifolds the d^H-cohomology has only a $\mathbb{Z}_2$-grading. For positive SKT manifolds, not only does the Laplacian preserve the spaces $\mathcal{W}_+^{k,l}$, but in fact there is an identity between the Laplacians for the operators d^H, δ_+^N and $\overline{\delta_+^N}$. We start with real Hodge theory and some operators of interest.

5.1 Differential operators, their adjoints and Laplacians

Given a generalized metric and orientation on a compact manifold M^n, we can form the Hodge star operator which gives us a positive definite inner product on forms:

$$\mathbb{G}(\varphi,\psi) = \int_M (\varphi, \star\psi)_{Ch} \qquad \text{for all } \varphi,\psi \in \Omega^\bullet(M;\mathbb{R}). \tag{5.1}$$

Two basic results about d^H are:

Lemma 5.1. If M^m is compact

$$\int_M (d^H\varphi,\psi)_{Ch} = (-1)^m \int_M (\varphi, d^H\psi)_{Ch}. \tag*{(Integration by parts)}$$

$$\mathbb{G}(d^H\varphi,\psi) = \mathbb{G}(\varphi, (-1)^m \star^{-1} d^H \star \psi) \tag*{(Formal adjoint)}$$

hence the formal adjoint of d^H is given by $d^{H*} = (-1)^{\frac{m(m+1)}{2}} \star d^H \star$.

The operators we will be mostly interested in are

$$\begin{aligned} \not{D}^H_+ &= \tfrac{1}{2}(d^H - (-1)^{\frac{m(m-1)}{2}} \star d^H \star) = \tfrac{1}{2}(d^H + (-1)^{m+1} d^{H*}) \\ \not{D}^H_- &= \tfrac{1}{2}(d^H + (-1)^{\frac{m(m-1)}{2}} \star d^H \star) = \tfrac{1}{2}(d^H + (-1)^{m} d^{H*}). \end{aligned} \tag{5.2}$$

The operators $\not{D}^H_\pm$ relate to the projections of d^H onto SD- and ASD- forms

$$d^H_\pm : \Omega^\bullet(M) \longrightarrow \Omega^\bullet_\pm(M),$$

$$d^H_\pm\varphi = \frac{1}{2}(d^H\varphi \mp i^{-\frac{m(m-1)}{2}}) \star d^H\varphi.$$

Lemma 5.2. Given $\varphi \in \Omega^\bullet(M)$ let $\varphi_\pm$ be its SD and ASD components, then

$$\not{D}^H_\pm\varphi = d^H_\mp\varphi_+ + d^H_\pm\varphi_-.$$

In particular we see that $\not{D}^H_-$ preserves the spaces $\Omega^\bullet_\pm(M)$ while $\not{D}^H_+$ maps them to each other.

Proof.

$$\begin{aligned} \not{D}^H_\pm\varphi &= \tfrac{1}{2}((d^H \mp (-1)^{\frac{m(m-1)}{2}} \star d^H \star)(\varphi_+ + \varphi_-) \\ &= \tfrac{1}{2}((d^H \pm (-1)^{\frac{m(m-1)}{2}} i^{\frac{m(m-1)}{2}} \star d^H)\varphi_+ + \tfrac{1}{2}((d^H \mp (-1)^{\frac{m(m-1)}{2}} i^{\frac{m(m-1)}{2}} \star d^H)\varphi_- \\ &= \tfrac{1}{2}((d^H \pm i^{3\frac{m(m-1)}{2}} \star d^H)\varphi_+ + \tfrac{1}{2}((d^H \mp i^{3\frac{m(m-1)}{2}} \star d^H)\varphi_- \\ &= \tfrac{1}{2}((d^H \pm i^{-\frac{m(m-1)}{2}} \star d^H)\varphi_+ + \tfrac{1}{2}((d^H \mp i^{-\frac{m(m-1)}{2}} \star d^H)\varphi_- \\ &= d^H_\mp\varphi_+ + d^H_\pm\varphi_-. \end{aligned}$$

□

We note that since $\not{D}^H_+$ swaps $\Omega^\bullet_+(M)$ with $\Omega^\bullet_-(M)$ it is a generalization of the signature operator. One could use this result to define an H-signature and an H-Euler characteritic as the indices of $\not{D}_\pm$. But since $\not{D}^H_+$ has the same symbol as $d+(-1)^{m+1}d^*$ and $\not{D}^H_-$ has the same symbol as $d+(-1)^m d^*$, the indices of these operators agree and hence they are just the usual signature and Euler characteristic.

Proposition 5.3. *Let $\triangle^H = d^H d^{H*} + d^{H*}d^H$ be the d^H-Laplacian. Then*

1. *$\star\triangle^H = \triangle^H\star$ and*
2. *$(-1)^{m+1}(\not{D}^H_+)^2 = (-1)^m(\not{D}^H_-)^2 = \frac{1}{4}\triangle^H$.*

Therefore $\triangle^H$ preserves the decomposition of forms into $\Omega^\bullet_\pm(M)$ and hence the d^H-cohomology of M splits as SD- *and* ASD-*cohomology: $H_{d^H}(M) = \mathcal{H}^H_+(M)\oplus\mathcal{H}^H_-(M)$.*

Proof. The first claim follows by simply expanding $d^{H*} = (-1)^{\frac{m(m+1)}{2}}\star d^H\star$:

$$\begin{aligned}\star\triangle^H &= (-1)^{\frac{m(m+1)}{2}}\star(d^H\star d^H\star+\star d^H\star d^H) = (-1)^{\frac{m(m+1)}{2}}(\star d^H\star d^H\star+\star^2 d^H\star d^H)\\ &= (-1)^{\frac{m(m+1)}{2}}(\star d^H\star d^H\star+d^H\star d^H\star^2) = \triangle^H\star.\end{aligned}$$

The second is just as straightforward:

$$(\not{D}^H_+)^2 = \tfrac{1}{4}(d^H+(-1)^{m+1}d^{H*})^2 = \tfrac{1}{4}(d^H)^2+(-1)^{m+1}(d^Hd^{H*}+d^{H*}d^H)+(d^{H*})^2 = \frac{(-1)^{m+1}}{4}\triangle^H.$$

□

5.2 Hodge theory on SKT manifolds

Theorem 5.4. *Let (M,H) be a compact manifold with 3-form, let $(\mathcal{G},\mathcal{I}_+,\mathcal{I}_-)$ be a generalized metric with a pair of almost Hermitian structures and $\mathcal{J}_1 = \mathcal{I}_+ + \mathcal{I}_-$. Then*

1. *If $\mathcal{I}_+$ (resp. $\mathcal{I}_-$) is a parallel positive (resp. negative) Hermitian structure, the d^H-Laplacian preserves the spaces $\mathcal{W}^{k,l}_+$ (resp. $\mathcal{W}^{k,l}_-$) and hence the d^H-cohomology of M inherits a corresponding $\mathbb{Z}\times\mathbb{Z}_2$-grading;*
2. *If $(\mathcal{G},\mathcal{J}_1)$ is a parallel bi-Hermitian structure, the d^H-Laplacian preserves the spaces $\mathcal{U}^{p,q}$ and hence the d^H-cohomology of M inherits a corresponding $\mathbb{Z}^2$-grading;*

Proof. The claim for positive and negative structures are analogous and together they imply the last claim, so it is enough to prove the first claim.

If (M,H) has a generalized Hermitian structure $(\mathcal{G},\mathcal{J}_1)$, then, since each diagonal of the $\mathcal{U}^{p,q}$ decomposition lies in either $\Omega^\bullet_+(M;\mathbb{C})$ or in $\Omega^\bullet_-(M;\mathbb{C})$, so Lemma 5.2 implies that $\not{D}^H_+$ and $\not{D}^H_-$ are decomposed into a sum of operators:

$$\begin{aligned}\not{D}^H_+ &= \delta_+ + N_1 + N_4 + N_+ + \overline{\delta}_+ + \overline{N_1} + \overline{N_4} + \overline{N_+};\\ \not{D}^H_- &= \delta_- + N_2 + N_3 + N_- + \overline{\delta}_- + \overline{N_2} + \overline{N_3} + \overline{N_-}.\end{aligned}\tag{5.3}$$

If $\mathcal{I}_+$ is parallel, then, according to Theorem 4.4, $N_2 = N_3 = 0$ and from (5.3) we have that $\not{D}^H_- = \delta_- + \overline{\delta}_- + N_- + \overline{N_-}$, which clearly preserves the $\mathcal{W}^k_+$ decomposition, therefore

$\triangle^H = 4(\not{D}^H_-)^2$ also preserves the $\mathcal{W}^k$ decomposition of forms. Since the Laplacian has even parity, it also preserves $\Omega^{ev}(M)$ and $\Omega^{od}(M)$, hence the Laplacian preserves the spaces $\mathcal{W}^{k,l}_+$. ☐

Corollary 5.5. Let M be a manifold with 3-form.

- An SKT structure on M induces a $\mathbb{Z} \times \mathbb{Z}_2$-grading on the d^H-cohomology;
- (Gualtieri [19]) A generalized Kähler structure induces a $\mathbb{Z}^2$-grading on the d^H-cohomology.

For each of the cases covered in the previous theorem, we denote by $H^{\bullet,\bullet}_{d^H}(M)$ the corresponding decomposition of the d^H-cohomology.

Corollary 5.6. Let (M^{2n}, H) be a compact manifold.

- If $(\mathcal{G}, \mathcal{I}_+)$ is a parallel positive (resp. negative) Hermitian structure for which $N_+ \not\equiv 0$ (resp. $N_- \not\equiv 0$), then $H^{\pm n,\bullet}_{d^H}(M) = \{0\}$.
- If $(\mathcal{G}, \mathcal{J}_1)$ is a parallel $U(n) \times U(n)$ structure on M for which $N_+ \not\equiv 0$ and $N_- \not\equiv 0$, then $H^{p,q}_{d^H}(M) = 0$ whenever $|p+q| = n$ or $|p-q| = n$.

Proof. Once again, the proof for positive and negative Hermitian structures is analogous and together they imply the claim for parallel bi-Hermitian structures, so we prove the claim only for positive structures. Assume that $H^{n,\bullet}(M)$ is nontrivial. Then there is a harmonic form $\rho \in \Gamma(\mathcal{W}^{n,\bullet}_+)$, in particular, ρ is d^H-closed, but the component of $d^H\rho$ in $\mathcal{W}^{n-3,\bullet}$ is given by $\overline{N_+}\rho$ and the action of $\wedge^\bullet V^{0,1}_+$ on $\mathcal{W}^n$ is faithful, hence $\overline{N_+}\rho = 0$ for $\rho \neq 0$ implies $\overline{N_+} = 0$, contradicting our initial hypothesis. ☐

Corollary 5.7. (Riemann bilinear relations) In a compact parallel positive Hermitian manifold

$$i^{-k}(a, \overline{a})_{Ch} > 0 \qquad \text{for all } a \in H^{k,l}_{d^H}(M)\backslash\{0\}.$$

Proof. Indeed, let α be the harmonic representative for the class $a \in H^{k,l}_{d^H}(M)\backslash\{0\}$. Since $\alpha \in \mathcal{W}^{k,l}_+$, $\star\alpha = i^k\alpha$ and hence $\overline{\star}\alpha = i^{-k}\overline{\alpha}$ and

$$i^{-k}(\alpha, \overline{\alpha})_{Ch} = (\alpha, \overline{\star}\alpha)_{Ch} > 0.$$

☐

Next we prove that the δ^N_+ Laplacian is a multiple of the d^H Laplacian in an SKT manifold. Firstly, we extend the real inner product on forms (5.1) to complex valued forms by requiring it to be Hermitian. If we let $\overline{\star}$ denote the operator given by $\overline{\star}\varphi = \star\overline{\varphi}$, we have

$$\mathbb{G}(\varphi, \psi) = (\varphi, \overline{\star}\psi) \qquad \text{for all } \varphi, \psi \in \Omega^\bullet(M;\mathbb{C}).$$

Proposition 5.8. *Let M^{2n} be a compact generalized almost Hermitian manifold and let δ denote any of the operators $\delta_+, \overline{\delta}_+, \delta_-, \overline{\delta}_-, N_\alpha$ or $\overline{N_\alpha}$, where $\alpha = 1, 2, 3, 4, \pm$. For $\varphi, \psi \in \Omega^\bullet(M;\mathbb{C})$ we have*

$$\int_M (\delta\varphi, \psi)_{Ch} = \int_M (\varphi, \delta\psi)_{Ch} \qquad \text{(Integration by parts)}$$

$$\delta^* = \pm\overline{\delta}, \qquad \text{(Formal adjoint)}$$

where the sign for the adjoint is positive if $\delta : \Omega_\pm(M;\mathbb{C}) \longrightarrow \Omega_\pm(M;\mathbb{C})$ *and negative otherwise.*

Proof. The proof that integration by parts holds is the same for all of these operators, so we consider only δ_+. It is enough to consider the case when $\varphi \in \mathcal{U}^{p,q}$ and hence $\delta_+\varphi \in \mathcal{U}^{p+1,q+1}$ and it pairs trivially with $\mathcal{U}^{k,l}$, unless $k = -p-1$ and $l = -q-1$. Hence we may assume that $\psi \in \mathcal{U}^{-p-1,-q-1}$ and compute

$$\int_M (\delta_+\varphi, \psi)_{Ch} = \int_M (d^H\varphi, \psi)_{Ch} = \int_M (\varphi, d^H\psi)_{Ch} = \int_M (\varphi, \delta_+\psi)_{Ch},$$

where in the first equality we have used that the remaining components of $d^H\varphi$ do not lie in $\mathcal{U}^{p+1,q+1}$, hence they pair trivially with ψ, in the second equality we integrated by parts and then reversed the argument.

For the formal adjoint, again taking $\delta = \delta_+$, $\varphi \in \mathcal{U}^{p,q}$ and $\psi \in \mathcal{U}^{p+1,q+1}$ we compute:

$$\begin{aligned}\mathbb{G}(\delta_+\varphi, \psi) &= \int_M (\delta_+\varphi, \overline{\star}\psi)_{Ch} = i^{-p-q-2}\int_M (\varphi, \delta_+\overline{\psi})_{Ch} \\ &= i^{-p-q-2}\int_M (\varphi, (-1)^n\overline{\star}\,\overline{\star}\delta_+\overline{\psi})_{Ch} = i^{-p-q-2}(-1)^n\int_M (\varphi, \overline{\star}\star\overline{\delta}_+\psi)_{Ch} \\ &= i^{-2p-2q-2}(-1)^n\int_M (\varphi, \overline{\star}\overline{\delta}_+\psi)_{Ch} = -\mathbb{G}(\varphi, \overline{\delta}_+\psi),\end{aligned}$$

where we have used several times that on $\mathcal{U}^{p,q}$ $\star$ is multiplication by i^{p+q} and in the last equality we used that $p+q = n \bmod 2$. ☐

Corollary 5.9. $(\delta_+^N)^* = -\overline{\delta_+^N}$.

Theorem 5.10. *In a positive SKT manifold,*

$$\triangle_{\delta_+^N} = \triangle_{\overline{\delta_+^N}} = \tfrac{1}{4}\triangle^H.$$

Proof. From $\not{D}_+^H = \delta_+^N + \overline{\delta_+^N}$ and Propositions 5.3 and 5.8 we have

$$\begin{aligned}\tfrac{1}{4}\triangle^H &= -(\not{D}_+^H)^2 = -(\delta_+^N + \overline{\delta_+^N})^2 = -(\delta_+^N)^2 - \delta_+^N\overline{\delta_+^N} - \overline{\delta_+^N}\delta_-^N - \overline{\delta_+^N}^2 \\ &= -\delta_+^N\overline{\delta_+^N} - \overline{\delta_+^N}\delta_+^N = \delta_+^N\delta_+^{N*} + \delta_+^{N*}\delta_+^N = \triangle_{\delta_+^N}\end{aligned}$$

☐

Corollary 5.11. The complex

$$\cdots \longrightarrow \mathcal{W}_+^{k-2} \xrightarrow{\delta_+^N} \mathcal{W}_+^k \xrightarrow{\delta_+^N} \mathcal{W}_+^{k+2} \xrightarrow{\delta_+^N} \cdots$$

is elliptic and hence so is the complex for δ_+.

Note that δ_-, obtained in the presence of a extension of the SKT structure to a generalized Hermitian structure, does not square to zero, hence the chain

$$\cdots \longrightarrow \mathcal{U}^{p-1,q-1} \xrightarrow{\delta_-} \mathcal{U}^{p,q} \xrightarrow{\delta_-} \mathcal{U}^{p+1,q+1} \xrightarrow{\delta_-} \cdots \tag{5.4}$$

is not a complex and its failure to be a complex is measured by $\delta_-^2 = -\{\delta_+, \overline{N}\}$.

5.3 A spectral sequence

The decomposition $\delta_+^N = \delta_+ + N$ obtained in the presence of a generalized Hermitian extension of a positive SKT structure is such that $\delta_+^2 = N^2 = \{\delta_+, N\} = 0$, hence this decomposition of the differential gives rise to a spectral sequence converging to the cohomology of δ_+^N. The first page of this sequence is just the δ_+-cohomology, $H^\bullet_{\delta_+}$. The next nontrivial page corresponds to the cohomology of the operator $N : H^\bullet_{\delta_+} \longrightarrow H^\bullet_{\delta_+}$ and the sequence goes on.

We can concretely compute the following pages of the sequence. Indeed, this is similar to the spectral sequence used to compute the cohomology of d^H from the usual de Rham cohomology [5, 1]. A given $\varphi^{p,q} \in \mathcal{U}^{p,q}$ gives rise to an element in the first page of the spectral sequence if it represents a nontrivial δ_+-cohomology class. Then $[\varphi^{p,q}]$ survives the next step if $[\varphi^{p,q}]$ is N-closed, that is $N \cdot [\varphi^{p,q}] = 0$, i.e., if there is $\varphi^{p-2,q+2} \in \mathcal{U}^{p-2,q+2}$ such that

$$N \cdot \varphi^{p,q} = \delta_+ \varphi^{p-2,q+2} \tag{5.5}$$

and it bounds if $[\varphi^{p,q}]$ is N-exact. If the class survived the second step, it will survive the third step if for some choice of $\varphi^{p-2,q+2}$ satisfying (5.5), $[\varphi^{p-2,q+2}]$ is N-closed and it will bound if it is N-exact, that is, we can find $\varphi^{p-4,q+4} \in \mathcal{U}^{p-4,q+4}$ such that

$$N \cdot \varphi^{p-2,q+2} = -\delta_+ \varphi^{p-4,q+4} \tag{5.6}$$

Notice that $N \cdot \varphi^{p-2,q+2}$ is just some kind of Massey product. Indeed, $N \cdot N = N^2 = 0 = \delta_+ 0$ and $N \cdot \varphi^{p,q} = \delta_+ \varphi^{p-2,q-2}$, so if we use the usual expression for Massey products of forms we have

$$\langle N, N, \varphi^{p,q} \rangle = N \varphi^{p-2,q+2}.$$

So the class of $\varphi^{p,q}$ continues to exist after the third step if the Massey product $\langle N, N, \varphi^{p,q} \rangle$ is defined and is the trivial δ_+-cohomology class.

The general step of the sequence corresponds to the Massey product $\langle \overbrace{N, N, \cdots, N}^{k}, \varphi^{p,q} \rangle$ and the class survived to this stage if there are choices of $\varphi^{p-jk,q+2j}$ for $j < k$, such that the previous products are defined and vanish. The class survives this stage if we can choose $\varphi^{p-2k,q+2k}$ with

$$\langle \overbrace{N, N, \cdots, N}^{k}, \varphi^{p,q} \rangle = -\delta_+ \varphi^{p-2k,q+2k}$$

and such that the Massey product

$$\langle \overbrace{N, N, \cdots, N}^{k}, \varphi^{p,q} \rangle$$

vanishes (in δ_+-cohomology).

Since on M^{2n} the bidegree of φ is bound by $\pm n$ we see that there are at most $n+1$ nontrivial steps in the sequence. The element $\varphi^{p,q}$ survives in E_∞ if all Massey products

$$\langle N, N, \cdots, N, \varphi^{p,q} \rangle$$

vanish in which case the forms φ^j above give rise to a δ_+^N-closed form: $\sum_{k=0}^{n+1} \varphi^{p-2k,q+2k}$.

With this argument we have proved



Figure 11: Graphic representation of the first four differentials of the spectral sequence relating δ_+- to δ_+^N-cohomology.

Theorem 5.12. *Let $(\mathcal{G},\mathcal{I}_+)$ be a positive SKT structure on a compact manifold with 3-form (M,H) and let $\mathcal{J}$ be a generalized complex structure extending $\mathcal{I}_+$. There is a spectral sequence converging to the δ_+^N-cohomology whose page E_2 is the δ_+ cohomology and whose differentials are Massey products*

$$d_k\varphi = \langle\overbrace{N,\cdots,N}^{k-1},\varphi\rangle.$$

This sequence degenerates after the page E_{n+1}.

Proposition 5.13. *Let $H^{k,l}_{d^H}(M)$ be the $\mathbb{Z}\times\mathbb{Z}_2$-graded decomposition of d^H-cohomology of a positive SKT manifold induced by the decomposition of forms into the spaces $W_+^{k,l}$. Also, let $H^{p,q}_{\delta_+}(M)$ be splitting of the cohomology of the operator δ_+ according to the $U^{p,q}$ decomposition. Then the Euler characteristic and the signature of M are given by*

$$\chi = \sum(-1)^l\dim(H^{n-2k,l}_{d^H}(M)) = \pm\sum(-1)^p\dim(H^{p,q}_{\delta_+}(M))$$

$$\sigma = \sum(-1)^k\dim(H^{n-2k,l}_{d^H}(M)) = \sum(-1)^{n-p-q}\dim(H^{p,q}_{\delta_+}(M))$$

Proof. As we argued earlier, the indices of $\not{D}^H_+$ and $\not{D}^H_-$ are just σ and χ, signature and Euler characteristic of M. Then the claims relating $H^{\bullet,\bullet}_{d^H}(M)$ with the χ and σ follow from the splitting of the d^H-cohomology obtained in 5.10.

As for δ_+, the proof is an iterated application of the fact that if

$$\{0\}\longrightarrow V_0\xrightarrow{d_0}V_1\xrightarrow{d_1}\cdots\xrightarrow{d_{n-1}}V_n\longrightarrow\{0\}.$$

is a complex, then the alternating sum of the dimensions is the same for the complex and its cohomology:

$$\sum(-1)^i\dim(V_i) = \sum(-1)^i\dim\left(\frac{\ker(d_i)}{\mathrm{Im}\,(d_{i-1})}\right),$$

and the fact that N maps $\Omega^{ev}(M) \longrightarrow \Omega^{od}(M)$ and $\Omega_+(M)$ into $\Omega_-(M)$ and vice versa, hence the corresponding alternating sum of the dimensions remains constant for all pages in the spectral sequence relating the δ_+-cohomology with the δ_+^N-cohomology. The sign to be used for the Euler characteristic in terms of $H_{\delta_+}(M)$ is positive if $\mathcal{U}^{0,n}$ is made of even forms and negative otherwise. ☐

5.4 Relation to Dolbeault cohomology

As we saw in Proposition 3.5, given a positive SKT structure $(\mathcal{G}, \mathcal{I})$ on M one can extend $\mathcal{I}_+$ to a generalized complex structure $\mathcal{J}_1$ of complex type. In this case, in the metric splitting of $\mathbb{T}M$, the structures $\mathcal{J}_1$ and $\mathcal{J}_2 = \mathcal{G}\mathcal{J}_1$ are given by

$$\mathcal{J}_1 = \begin{pmatrix} I & 0 \\ 0 & -I^* \end{pmatrix}, \qquad \mathcal{J}_2 = \begin{pmatrix} 0 & -\omega^{-1} \\ \omega & 0 \end{pmatrix},$$

where $\omega = g \circ I$ and for such generalized Hermitian structure Proposition 1.9 gives an automorphism of the space of forms which relates the $U^{p,q}$ decomposition of forms with the usual $\wedge^{p,q}T^*$ decomposition of forms determined by the complex structure I.

In this section we relate these two decomposition of forms and corresponding cohomologies. Precisely, the effect of applying the automorphism Ψ from (1.7) to $\wedge^\bullet T^*_{\mathbb{C}}M$ on d^H is simply to conjugate it by Ψ, so we get a new operator

$$\hat{d^H} = \Psi^{-1} \circ d^H \circ \Psi.$$

Since d^H splits according to the $U^{p,q}$ into six operators, the same is true for $\hat{d^H}$ and we can define

$$\begin{aligned} &\hat{\delta}_+ : \Omega^{p,q} \longrightarrow \Omega^{p-1,q}; && \hat{\overline{\delta}}_+ : \Omega^{p,q} \longrightarrow \Omega^{p+1,q}; \\ &\hat{\delta}_- : \Omega^{p,q} \longrightarrow \Omega^{p,q+1}; && \hat{\overline{\delta}}_- : \Omega^{p,q} \longrightarrow \Omega^{p,q+1}; \\ &\hat{N} : \Omega^{p,q} \longrightarrow \Omega^{p-1,q-2} && \hat{\overline{N}} : \Omega^{p,q} \longrightarrow \Omega^{p+1,q+2}; \end{aligned}$$

$$\hat{d^H} = \hat{\delta}_+ + \hat{\overline{\delta}}_+ + \hat{\delta}_- + \hat{\overline{\delta}}_- + \hat{N} + \hat{\overline{N}}.$$

Proposition 5.14. *Let $(\mathcal{G}, \mathcal{I}_+)$ be a positive SKT structure on a manifold with 3-form (M, H), let $\mathcal{J}$ be the generalized complex extension of $\mathcal{I}_+$ given by Proposition 3.5. Let*

$$\partial^{\omega^{-1}} = \{\partial, \omega^{-1}\} \qquad \textit{and} \qquad \overline{\partial}^{\omega^{-1}} = \{\overline{\partial}, \omega^{-1}\},$$

let ζ_i be the component of $e_*^{\frac{i}{2}\omega^{-1}}\overline{\partial}\omega$ lying in $\wedge^i T \otimes \wedge^{3-i}T^*$ and let χ be the $(1,2)$-component of $[\omega^{-1}, \omega^{-1}]_{SN}$, where $[\cdot,\cdot]_{SN}$ is the Schouten–Nijenhuis bracket. Then the following hold

$$\begin{aligned} &\hat{\delta}_+ = \tfrac{i}{2}\overline{\partial}^{\omega^{-1}} + 2i\zeta_2 && \hat{\overline{\delta}}_+ = \partial \\ &\hat{\delta}_- = \overline{\partial} + 2i\zeta_1 && \hat{\overline{\delta}}_- = \tfrac{i}{2}\partial^{\omega^{-1}} \\ &\hat{N} = \tfrac{1}{8}\chi && \hat{\overline{N}} = 2i\overline{\partial}\omega. \end{aligned} \tag{5.7}$$

Proof. The proof is a direct computation of the operator $\hat{d^H}$:

$$\begin{aligned} \hat{d^H} &= e^{-\frac{i}{2}\omega^{-1}} e^{-i\omega}(d + H\wedge)e^{i\omega}e^{\frac{i}{2}\omega^{-1}} \\ &= e^{-\frac{i}{2}\omega^{-1}} e^{-i\omega}e^{i\omega}(d + (H + id\omega)\wedge)e^{\frac{i}{2}\omega^{-1}} \\ &= e^{-\frac{i}{2}\omega^{-1}} de^{\frac{i}{2}\omega^{-1}} + 2ie^{-\frac{i}{2}\omega^{-1}}(\overline{\partial}\omega\wedge)e^{\frac{i}{2}\omega^{-1}} \end{aligned} \tag{5.8}$$

We compute separately each of the two operators above making up $\hat{d^H}$:

$$e^{-\frac{i}{2}\omega^{-1}} de^{\frac{i}{2}\omega^{-1}} = d + [d, \tfrac{i}{2}\omega^{-1}] + \tfrac{1}{2!}[[d, \tfrac{i}{2}\omega^{-1}], \tfrac{i}{2}\omega^{-1}] + \tfrac{1}{3!}[[[d, \tfrac{i}{2}\omega^{-1}], \tfrac{i}{2}\omega^{-1}], \tfrac{i}{2}\omega^{-1}] + \cdots$$

The first term is just $d = \partial + \overline{\partial}$, while the second term is a version of the symplectic adjoint of d introduced by Koszul [26], but now obtained in a nonintegrable setting, $\delta^{\omega^{-1}} = d \circ \omega^{-1} - \omega^{-1} \circ d$. Since ω is not closed, $\delta^{\omega^{-1}}$ does not square to zero. Integrability of the complex structure gives $d = \partial + \overline{\partial}$ and recalling that ω^{-1} is of type $(1, 1)$ we have

$$\delta^{\omega^{-1}} = \partial^{\omega^{-1}} + \overline{\partial}^{\omega^{-1}}; \partial^{\omega^{-1}} : \Omega^{p,q} \longrightarrow \Omega^{p,q-1}; \qquad \overline{\partial}^{\omega^{-1}} : \Omega^{p,q} \longrightarrow \Omega^{p-1,q}.$$

Since ω^{-1} is even, the third term in the series coincides with a multiple of the expression for the derived bracket of ω^{-1} with itself, that is, the Schouten–Nijenhuis bracket:

$$[[d, \tfrac{i}{2}\omega^{-1}], \tfrac{i}{2}\omega^{-1}] = -\{\{\tfrac{i}{2}\omega^{-1}, d\}, \tfrac{i}{2}\omega^{-1}\} = \tfrac{1}{4}[\omega^{-1}, \omega^{-1}]_{SN}.$$

Since ω^{-1} is of type $(1, 1)$, integrability of the complex structure implies that this term this is a 3-vector lies in $\wedge^{2,1}T \oplus \wedge^{1,2}T$, so this terms also decomposes:

$$[\omega^{-1}, \omega^{-1}]_{SN} = \chi + \overline{\chi}, \qquad \chi \in \wedge^{1,2}TM.$$

The fourth term is the commutator $[[\omega^{-1}, \omega^{-1}]_{SN}, \omega^{-1}]$ which vanishes since ω^{-1} is a bivector and hence so do the remaining terms in the series. So we have established that

$$e^{-\frac{i}{2}\omega^{-1}} de^{-\frac{i}{2}\omega^{-1}} = \partial + \overline{\partial} + \tfrac{i}{2}\partial^{\omega^{-1}} + \tfrac{i}{2}\overline{\partial}^{\omega^{-1}} + \tfrac{1}{8}(\chi + \overline{\chi}) \tag{5.9}$$

Next we compute the second summand in (5.8):

$$\begin{aligned} 2ie^{-\frac{i}{2}\omega^{-1}} \circ \overline{\partial}\omega \circ e^{\frac{i}{2}\omega^{-1}} &= 2ie^{-\frac{i}{2}\omega^{-1}} \circ (e_*^{\frac{i}{2}\omega^{-1}} e_*^{-\frac{i}{2}\omega^{-1}}\overline{\partial}\omega) \circ e^{\frac{i}{2}\omega^{-1}} \\ &= 2ie^{-\frac{i}{2}\omega^{-1}} \circ e^{\frac{i}{2}\omega^{-1}} \circ (e_*^{-\frac{i}{2}\omega^{-1}}\overline{\partial}\omega) \\ &= 2ie_*^{-\frac{i}{2}\omega^{-1}}\overline{\partial}\omega, \end{aligned}$$

where we have used (1.1) in the second equality. The element $e_*^{-\frac{i}{2}\omega^{-1}}\overline{\partial}\omega \in \wedge^3\mathbb{T}_{\mathbb{C}}M$ has six components:

$$e_*^{-\frac{i}{2}\omega^{-1}}\overline{\partial}\omega \in (\wedge^{2,1}T^*)\oplus(T^{1,0}\otimes\wedge^{1,1}T^*)\oplus(T^{0,1}\otimes\wedge^{0,2}T^*)\oplus(\wedge^{2,0}T\otimes T^{*1,0})\oplus(\wedge^{1,1}T\otimes T^{*0,1})\oplus(\wedge^{1,2}T)$$

Since, for a 1-form ξ, $e_*^{-\frac{i}{2}\omega^{-1}}\xi = \frac{i}{2}\omega^{-1}(\xi) + \xi$, we see that the 3-form component is $\overline{\partial}\omega$ and that the 3-vector component is $-\frac{i}{8}\omega^{-1}\overline{\partial}\omega$. We let ζ_1 be the component of

$e_*^{-\frac{i}{2}\omega^{-1}}\overline{\partial}\omega$ in $T\otimes\wedge^2T^*$ and ζ_2 be the component in $\wedge^2T\otimes T^*$, so that $\zeta_1:\wedge^{p,q}\longrightarrow\wedge^{p,q+1}$, $\zeta_2:\wedge^{p,q}\longrightarrow\wedge^{p-1,q}$, and

$$2ie^{-\frac{i}{2}\omega^{-1}}\circ\overline{\partial}\omega\circ e^{\frac{i}{2}\omega^{-1}}=2i\overline{\partial}\omega+2i\zeta_1+2i\zeta_2+\frac{1}{4}\omega^{-1}(\overline{\partial}\omega)$$

Putting this together with (5.9) and the fact that $\hat{d^H}$ does not have a component mapping $\Omega^{p,q}(M)$ to $\Omega^{p-2,q-1}(M)$ we conclude that $\overline{\chi}=2\omega^{-1}(\overline{\partial}\omega)$ and obtain (5.7). $\square$

Next, we let $\partial^{i\overline{\partial}\omega}$ be the operator $\partial+i\overline{\partial}\omega\wedge$. Then we observe that even though $\partial^{i\overline{\partial}\omega}$ does not preserve the degree of a form, it preserves the holomorphic degree, i.e.,

$$\partial^{i\overline{\partial}\omega}:\Omega^{p,\bullet}(M)\longrightarrow\Omega^{p+1,\bullet}(M),$$

so its cohomology has a natural $\mathbb{Z}$-grading. For an operator δ with $\delta^2=0$ we denote its cohomology by $H_\delta(M)$.

Corollary 5.15. Let $(\mathcal{G},\mathcal{I}_+)$ be a positive SKT structure on a compact manifold, let $\mathcal{J}$ be the generalized complex extension from Proposition 3.5 and let (g,I) be the corresponding Hermitian structure with Hermitian form $\omega=g\circ I$. Then

$$H^{p,q}_{\partial}(M)\cong H^{q-p,n-p-q}_{\overline{\delta_+}}(M),$$

and

$$H^{p}_{\partial^{i\overline{\partial}\omega}}(M)\cong H^{n-2p}_{\overline{\delta^N_+}}(M)\cong H^{n-2p}_{d^H}(M).$$

Proof. Indeed, Ψ puts $\Omega^{p,q}(M)$ in correspondence with $\mathcal{U}^{q-p,n-p-q}$ and $\Omega^{p,\bullet}(M)$ with $\mathcal{W}^{n-2p}$ and according to Proposition 5.14, $\Psi\circ\partial=\overline{\delta_+}\circ\Psi$ and $\Psi\circ\partial^{2i\omega}=\overline{\delta^N_+}\circ\Psi$. Therefore the $H^{p,q}_{\partial}(M)\cong H^{q-p,n-p-q}_{\overline{\delta_+}}(M)$ and $H^{p}_{\partial^{2i\overline{\partial}\omega}}(M)\cong H^{n-2p}_{\overline{\delta^N_+}}(M)$. Of course the cohomology of $\partial^{2i\overline{\partial}\omega}$ is the same as the cohomology if $\partial^{i\overline{\partial}\omega}$, as the automorphism

$$m:\wedge^{\bullet,\bullet}T^*M\longrightarrow m:\wedge^{\bullet,\bullet}T^*M\qquad m(\alpha)=2^{\frac{q}{2}}\alpha\qquad\text{for all }\alpha\in\wedge^{p,q}T^*M,$$

relates them. $\square$

Interestingly we have that $d^H=\partial^{i\overline{\partial}\omega}+\overline{\partial}^{-i\partial\omega}$ and the operators $\partial^{i\overline{\partial}\omega}$ and $\overline{\partial}^{-i\partial\omega}$ both square to zero (due to the SKT condition $\partial\overline{\partial}\omega=0$) and graded commute. Yet, they are not well behaved with respect to the $\mathbb{Z}^2$-bigrading of forms and only on a $\mathbb{Z}_2^2$-grading do they behave nicely, as there we have

$$\partial^{i\overline{\partial}\omega}:\Omega^{p,q}(M)\longrightarrow\Omega^{p+1,q}(M)\qquad\overline{\partial}^{-i\partial\omega}:\Omega^{p,q}(M)\longrightarrow\Omega^{p,q+1}(M),\qquad(p,q)\in\mathbb{Z}_2^2.$$

There is a standard trick to recover one $\mathbb{Z}$-grading, say the 'p'-grading. We introduce a new formal variable, β, alter the operators $\partial^{i\overline{\partial}\omega}$ and $\overline{\partial}^{-i\partial\omega}$ so that they act on $E_1=\Omega^{p,q}(M)\otimes\langle\beta\rangle$:

$$\partial^{i\overline{\partial}\omega}(\alpha\beta^k)=(\partial^{i\overline{\partial}\omega}\alpha)\beta^k\qquad\overline{\partial}^{-i\partial\omega}(\alpha\beta^k)=(\overline{\partial}\alpha)\beta^k-i\partial\omega\wedge\alpha\beta^{k+1}$$

and introduce a $\mathbb{Z}\times\mathbb{Z}_2$-grading on E_1 by declaring that for $\Omega^{p,q}(M)\otimes\beta^k = E_1^{p-2k,q}$, $(p-2k,q)\in\mathbb{Z}\times\mathbb{Z}_2$. I.e., β has degree $(-2,0)$. With this arrangement

$$\partial^{i\overline{\partial}\omega} : E_1^{p,q}(M) \longrightarrow E_1^{p+1,q}(M) \qquad \overline{\partial}^{-i\partial\omega} : E_1^{p,q}(M) \longrightarrow E_1^{p,q+1}(M), \qquad (p,q)\in\mathbb{Z}\times\mathbb{Z}_2.$$

In this setting, one can produce a $\mathbb{Z}\times\mathbb{Z}_2$-graded spectral sequence whose second page is $H^\bullet_{\partial^{i\overline{\partial}\omega}}(M)\otimes\langle\beta\rangle$ and the last is $H^\bullet_{d^H}(M)\otimes\langle\beta\rangle$. Since these are isomorphic, by Corollary 5.15, this sequence degenerates at the second page.

Theorem 5.16. (SKT Frölicher spectral sequence) *In an SKT manifold the decomposition* $d^H = (\partial + i\overline{\partial}\omega) + (\overline{\partial} - i\partial\omega)$ *gives rise to a spectral sequence which degenerates at the second page.*

5.5 Alternative definition

One shortcoming of SKT structures compared the Kähler structures is that there is no analogue of the d^c-operator. Instead, there are two operators, neither of which fully works as the complex d^c. Indeed, given a positive SKT structure $(\mathcal{G},\mathcal{I})$, we can define $\delta^{\mathcal{I}}$ and $d^{\mathcal{I}}$ as

$$\begin{aligned}\delta^{\mathcal{I}} &= \{d^H,\mathcal{I}\} = -i(\delta_+^N - \overline{\delta_+^N})\\ d^{\mathcal{I}} &= e^{\frac{-\pi\mathcal{I}}{2}} d^H e^{\frac{\pi\mathcal{I}}{2}} = -i(\delta_+^N - \overline{\delta_+^N}) + \not{\delta}_-;\end{aligned} \tag{5.10}$$

Lemma 5.17. The following hold:

$$\delta^{\mathcal{I}*} = -\overline{\delta^{\mathcal{I}}};$$

$$(\delta^{\mathcal{I}})^2 = -\tfrac{1}{4}\triangle^H \qquad \{\delta^{\mathcal{I}}, d^H\} = 0.$$

$$(d^{\mathcal{I}})^2 = 0 \qquad \{d^{\mathcal{I}}, d^H\} = \tfrac{1}{2}\triangle^H;$$

$$(d^{\mathcal{I}^*})^2 = 0 \qquad \{d^{\mathcal{I}*}, d^H\} = \tfrac{1}{2}\triangle^H;$$

Proof. The claim $\delta^{\mathcal{I}*} = -\overline{\delta^{\mathcal{I}}}$ follows from 5.9. Using Corollary 3.4, Corollary 5.9 and Theorem 5.10 we have

$$(\delta^{\mathcal{I}})^2 = -(\delta_+^N - \overline{\delta_+^N})^2 = \delta_+^N\overline{\delta_+^N} + \overline{\delta_+^N}\delta_+^N = -(\delta_+^N\delta_+^{N*} + \delta_+^{N*}\delta_+^N) = -\tfrac{1}{4}\triangle^H.$$

Using that the graded commutator satisfies the graded Jacobi identity we have

$$\{d^H,\delta^{\mathcal{I}}\} = \{d^H,\{d^H,\mathcal{I}\}\} = \{\{d^H,d^H\},\mathcal{I}\} - \{d^H,\{d^H,\mathcal{I}\}\} = -\{d^H,\delta^{\mathcal{I}}\},$$

and therefore $\{d^H,\delta^{\mathcal{I}}\} = 0$.

The conditions $(d^{\mathcal{I}})^2 = (d^{\mathcal{I}*})^2 = 0$ are obvious and for the last equalities we use that $d^{\mathcal{I}} = \delta^{\mathcal{I}} + \not{D}_-^H$ and $d^{\mathcal{I}*} = -\delta^{\mathcal{I}} + \not{D}_-^H$, so, for example

$$\{d^{\mathcal{I}}, d^H\} = \{\delta^{\mathcal{I}} + \not{D}_-^H, d^H\} = \{\not{D}_-^H, d^H\} = \{\not{D}_-^H, \not{D}_-^H\} + \{\not{D}_-^H, \not{D}_+^H\} = \tfrac{1}{2}\triangle^H + 0.$$

□

Since $(d^{\mathcal{I}})^2 = 0$, one can also study its cohomology. Not surprisingly, we have the following corollary to Theorem 5.10

Corollary 5.18. Let $\triangle_{d^{\mathcal{I}}}$ be the Laplacian of $d^{\mathcal{I}}$. Then

$$\triangle_{d^{\mathcal{I}}} = \triangle^H.$$

Proof. Indeed, since $d^{\mathcal{I}} = \delta^{\mathcal{I}} + \not{D}^H_-$, then $d^{\mathcal{I}*} = -\overline{\delta^{\mathcal{I}}} + \not{D}^H_-$ and we have

$$\begin{aligned}\triangle_{d^{\mathcal{I}}} &= \{d^{\mathcal{I}}, d^{\mathcal{I}*}\}\\ &= \{\delta^{\mathcal{I}} + \not{D}^H_-, -\overline{\delta^{\mathcal{I}}} + \not{D}^H_-\}\\ &= -\{\delta^{\mathcal{I}}, \overline{\delta^{\mathcal{I}}}\} + \{\not{D}^H_-, \not{D}^H_-\}\\ &= \triangle^H,\end{aligned}$$

where in the fourth equality we used Corollary 3.4, in the fifth we used that $\overline{\delta^N_+} = -\delta^{N*}_+$ and Proposition 5.3 and in the last equality we used Theorem 5.10 ☐

In light of Lemma 5.17, there seems to be no analogue of the dd^c-lemma in the SKT setting as we do not have enough operators satisfying the correct relations. Yet the operator $\delta^{\mathcal{I}}$ can be useful in a different context, as we explain next. In [17], Grabowski showed that a generalized complex structure is equivalent to a tensor $\mathcal{J} \in \Gamma(\wedge^2\mathbb{T}M) = \Gamma(\mathfrak{spin}(\mathbb{T}M))$ acting on forms via the spin action on spinors and for which the following relation holds

$$\{\{d^H, \mathcal{J}\}, \mathcal{J}\} = -d^H.$$

One can see where this comes from: for a generalized complex structure we define $d^{\mathcal{J}} = \{d^H, \mathcal{J}\} = -i(\partial - \overline{\partial})$ and hence $\{d^{\mathcal{J}}, \mathcal{J}\} = (-i)^2(\partial + \overline{\partial}) = -d^H$. So this equation includes simultaneously the fact that $\mathcal{J}^2 = -\mathrm{Id}$ and the compatibility between $\mathcal{J}$ and d^H, i.e., the integrability condition. Further, it is phrased in a way which allows one to work with generalized complex structures in the framework of super geometry.

In our setting we can find a similar condition:

Theorem 5.19. *Let (M^{2n}, H) be an oriented manifold with 3-form and $\mathcal{G}$ be a generalized metric on M. A tensor $\widetilde{\mathcal{I}} \in \Gamma(\wedge^2 V_+) \subset \Gamma(\wedge^2\mathbb{T}M)$ is an SKT structure if and only if*

$$\{\{d^H, \widetilde{\mathcal{I}}\}, \widetilde{\mathcal{I}}\} = -\not{D}^H_+, \tag{5.11}$$

Proof. One implication is easy. If $\mathcal{I}$ is an SKT structure. Then $\{d^H, \mathcal{I}\} = -i(\delta^N_+ - \overline{\delta^N_+})$ and $\{\{d^H, \mathcal{I}\}, \mathcal{I}\} = (-i)^2(\delta^N_+ + \overline{\delta^N_+}) = -\not{D}^H_+$.

The converse requires a little more effort. We start with a pointwise argument. Since $\widetilde{\mathcal{I}} \in \Gamma(\wedge^2\mathbb{T}M) = \Gamma(\mathfrak{spin}(\mathbb{T}M))$, it acts on forms, thought of as spinors, in a skew symmetric way (with respect to the Chevalley pairing). Since $\widetilde{\mathcal{I}} \in \Gamma(\wedge^2 V_+)$, Lemma 1.5 implies that $\star\widetilde{\mathcal{I}} = \widetilde{\mathcal{I}}\star$ which means, firstly, that $\widetilde{\mathcal{I}}$ preserves the spaces $\Omega^\bullet_\pm(M)$ and secondly that $\widetilde{\mathcal{I}}$ is skew-symmetric with respect to the metric on forms induced by the Hodge star (1.2). Therefore, the Clifford action of $\widetilde{\mathcal{I}}$ on forms is diagonalizable and its eigenvalues are purelly imaginary numbers. We denote the $i\lambda$-eigenspace of $\widetilde{\mathcal{I}}$ by $W^{2\lambda}$.

Similarly, there is a bundle endomorphism of $\mathbb{T}M$ associated to $\widetilde{\mathcal{I}}$ given by

$$\mathcal{I}v = \{\widetilde{\mathcal{I}}, v\}, \quad v \in \mathbb{T}M.$$

using again that $\widetilde{\mathcal{I}} \in \Gamma(\wedge^2 V_+)$, we conclude that if $v \in V_-$, then $\mathcal{I}v = \{\widetilde{\mathcal{I}}, v\} = 0$, and if $v \in V_+$ then $\mathcal{I}v \in V_+$. Further $\mathcal{I}$, as an endomorphism of V_+, is skew symmetric with respect to the natural pairing which is positive definite on V_+. Hence $\mathcal{I}$ is also diagonalizable and has purely imaginary eigenvalues.

If v is an $i\mu$-eigenvector of $\mathcal{I}$ and $\rho \in W^{2\lambda}$, then

$$\widetilde{\mathcal{I}}(v \cdot \rho) = \{\widetilde{\mathcal{I}}, v\} \cdot \rho + v \cdot (\widetilde{\mathcal{I}}\rho) = (\mathcal{I}v) \cdot \rho + v \cdot (\widetilde{\mathcal{I}}\rho) = (\mu + \lambda) i v \cdot \rho, \tag{5.12}$$

that is, the eigenvectors of $\mathcal{I}$ map eigenspaces of $\widetilde{\mathcal{I}}$ to each other.

Next we use a local argument. Notice that the eigenvalues of $\widetilde{\mathcal{I}}$ and of $\mathcal{I}$ may vary from point to point and hence the eigenspaces are not necessarily vector bundles, as there may be a locus in which generically distinct eigenvalues coincide. So, in what follows, we only consider an open set where the eigenspaces are actually vector bundles and denote by $\mathcal{W}^{2\lambda}$ the space of sections of the eigenbunde corresponding to eigenfunction $i\lambda$.

Let $\rho \in \mathcal{W}^{2\lambda}$ be, say, a (locally) nonvanishing SD-form and decompose $d^H\rho$ into components:

$$d^H\rho = \sum_\mu \rho_\mu^+ + \rho_\mu^-,$$

where $\rho_\mu^\pm \in \mathcal{W}^{2\mu} \cap \Omega^\bullet_\pm(M)$, so that

$$\not{D}_+^H \rho = \sum_\mu \rho_\mu^-, \qquad \not{D}_-^H = \sum_\mu \rho_\mu^+. \tag{5.13}$$

Then

$$\begin{aligned}
\{\{d^H, \widetilde{\mathcal{I}}\}, \widetilde{\mathcal{I}}\}\rho &= (\widetilde{\mathcal{I}}^2 d^H - 2\widetilde{\mathcal{I}} d^H \widetilde{\mathcal{I}} + d^H \widetilde{\mathcal{I}}^2)\rho \\
&= \widetilde{\mathcal{I}}^2 d^H\rho - 2i\widetilde{\mathcal{I}} d^H(\lambda\rho) - d^H(\lambda^2\rho) \\
&= \widetilde{\mathcal{I}}^2 d^H\rho - 2i\widetilde{\mathcal{I}}(d\lambda \wedge \rho) - 2i\lambda\widetilde{\mathcal{I}} d^H\rho - 2\lambda d\lambda \wedge \rho - \lambda^2 d^H\rho \\
&= \widetilde{\mathcal{I}}^2 d^H\rho - 2i\lambda\widetilde{\mathcal{I}} d^H\rho - \lambda^2 d^H\rho - 2i(\mathcal{I}d\lambda) \cdot \rho - 2i(d\lambda) \wedge \widetilde{\mathcal{I}}\rho - 2\lambda d\lambda \wedge \rho \\
&= \sum_\mu (-\mu^2 + 2\mu\lambda - \lambda^2)(\rho_\mu^+ + \rho_\mu^-) - 2i(\mathcal{I}d\lambda) \cdot \rho \\
&= -\sum_\mu (\mu - \lambda)^2(\rho_\mu^+ + \rho_\mu^-) - 2i(\mathcal{I}d\lambda) \cdot \rho
\end{aligned}$$

and since $\mathcal{I} : \mathbb{T}M \longrightarrow V_+$, $2i(\mathcal{I}d\lambda) \cdot \rho$ is an ASD-form, hence the SD and ASD parts of the equation above are

$$(\{\{d^H, \widetilde{\mathcal{I}}\}, \widetilde{\mathcal{I}}\}\rho)_+ = -\sum_\mu (\mu - \lambda)^2 \rho_\mu^+$$

$$(\{\{d^H, \widetilde{\mathcal{I}}\}, \widetilde{\mathcal{I}}\}\rho)_- = -2i(\mathcal{I}d\lambda) \cdot \rho - \sum_\mu (\mu - \lambda)^2 \rho_\mu^-.$$

Hence, condition (5.11) is equivalent to

$$\sum_{\mu}(\mu-\lambda)^2\rho^+_\mu = 0, \tag{5.14}$$

$$2i(\mathcal{I}d\lambda)\cdot\rho + \sum_{\mu}(\mu-\lambda)^2\rho^-_\mu = \sum_{\mu}\rho^-_\mu. \tag{5.15}$$

the first equation implies that the only value of μ for which ρ^+_μ may be non zero is $\mu=\lambda$ and hence it follows from (5.13) that $\not{D}^H_-\rho = \rho^+_\lambda$, that is

$$\not{D}^H_- : \mathcal{W}^\lambda \longrightarrow \mathcal{W}^\lambda. \tag{5.16}$$

Equation (5.15) has more implications. Firstly, if we denote the component of $d\lambda$ in the $i\nu$-eigenspace of $\mathcal{I}$ by $(d\lambda)_\nu$, then for $\nu \neq \pm 1$, taking projection of (5.15) onto $\mathcal{W}^{2\lambda+2\nu}$ we obtain the identity

$$-2\nu(d\lambda)_\nu\cdot\rho + \nu^2\rho^-_{\lambda+\nu} = \rho^-_{\lambda+\nu}$$

and for $\nu \neq \pm 1$ we get

$$\rho^-_{\lambda+\nu} = \frac{2\nu(d\lambda)_\nu}{\nu^2-1}\cdot\rho.$$

In particular, we see that for $\nu \neq \pm 1$ the component of $d^H\rho$ in $\mathcal{W}^{2\lambda+2\nu}$ depends tensorially on $\rho \in \mathcal{W}^{2\lambda}$.

On the other hand, since $d^H(f\rho) = df\rho + fd^H\rho$, if we project this onto $\mathcal{W}^{2\lambda+2\mu}$ we get

$$\frac{2\nu(d\lambda)_\nu}{\nu^2-1}\cdot f\rho = (d^H(f\rho))_{\lambda+\mu} = (df)_\mu\cdot\rho + f(d^H\rho)_{\lambda+\mu} = (df)_\nu\cdot\rho + f\frac{2\nu(d\lambda)_\nu}{\nu^2-1}\cdot\rho.$$

That is $(df)_\nu$ must annihilate $\rho \in \mathcal{W}^{2\lambda}\cap\Omega_+(M)$ for every $\nu \neq \pm 1$. Since this holds for every λ, and the same argument can be used when ρ is an ASD form, we conclude that $(df)_\nu$ annihilates all forms and hence must vanish if $\nu \neq \pm 1$. Therefore $\mathcal{I}|_{V_+}$ has only $\pm i$ as eigenvalues. This means that the eigenspaces of $\mathcal{I}$ actually form a subbundle of $\mathbb{V}_+\otimes\mathbb{C}$ and $\mathcal{I}$ is a complex structure on V_+. Since the action of $\mathrm{Clif}(\mathbb{T}_\mathbb{C}M)$ on forms is transitive and the only eigenvalues of $\mathcal{I}$ are 0 and $\pm i$ we conclude from (5.12) that the the eigenvalues of $\widetilde{\mathcal{I}}$ differ by imaginary integers and since $\widetilde{\mathcal{I}}$ is real, complex conjugation swaps $\mathcal{W}^{2\lambda}$ with $W^{-2\lambda}$, hence the eigenvalues of $\widetilde{\mathcal{I}}$ must be either all of the form ik or of the form $i(k+\frac{1}{2})$ for $k\in\mathbb{Z}$. Either way, the eigenvalues of $\widetilde{\mathcal{I}}$ are constant, $2\lambda = k$ for some $k\in\mathbb{Z}$ and (5.15) becomes

$$\not{D}^H_+\rho = \rho^-_{k+1} + \rho^-_{k-1} \in \mathcal{W}^{2k+2}\oplus\mathcal{W}^{2k-2}.$$

Adding this to (5.16) we obtain

$$d^H : \mathcal{W}^k \longrightarrow \mathcal{W}^{k+2} + \oplus\mathcal{W}^k \oplus \mathcal{W}^{k-2},$$

showing that $\mathcal{I}$ is an integrable SKT structure due to Theorem 3.3. □

Corollary 5.20. A generalized Kähler structure is a generalized metric $\mathcal{G}$ and two tensors $\mathcal{I}_\pm \in \Gamma(\wedge^2 V_\pm)$ such that

$$\{\{d^H,\mathcal{I}_\pm\},\mathcal{I}_\pm\} = -\not{D}^H_\pm, \tag{5.17}$$

6 Instantons over complex surfaces

Next we use the techniques developed in this section to provide an alternative description of the SKT structure on the moduli space of instantons of a bundle over a complex surface. The tool used to describe this structure are extended actions as introduced in [3] and the SKT reduction theorem as presented in [7]. The argument presented here follows closely the one from [4], so we will spare details and refer to that paper for further reading.

Let $(M,[g],I)$ be a compact complex surface with a conformal Hermitian structure. By a result of Gauduchon [15], there is a representative g of the conformal class which makes (M,g,I) into an SKT manifold, i.e., the corresponding Hermitian form ω satisfies $dd^c\omega = 0$. We let $H = d^c\omega$ and consider $\mathbb{T}M$ endowed with the H-Courant bracket, so that the metric

$$\mathcal{G} = \begin{pmatrix} 0 & g^{-1} \\ g & 0 \end{pmatrix}$$

and the complex structure on V_+ induced by I via the isomorphism $\pi : V_+ \longrightarrow TM$ are a positive SKT structure on $\mathbb{T}M$.

Given a bundle E over M with a compact Lie group G as structure group and Lie algebra $\mathfrak{g}$ we let $\mathbb{A}$ be the space of all $\mathfrak{g}$-connections on E endowed with the trivial 3-form, so that $\mathbb{A}$ is an affine space isomorphic to space of 1-forms on M with values in the adjoint bundle $\mathfrak{g}_E$, $\Omega^1(M;\mathfrak{g}_E)$. Hence at any connection D we have $T_D\mathbb{A} = \Omega^1(M;\mathfrak{g}_E)$ and, letting κ be a bi-invariant metric on G, we can use κ to identify $T^*_D\mathbb{A} = \Omega^3(M;\mathfrak{g}_E)$. Indeed, for $X \in \Omega^1(M;\mathfrak{g}_E)$ and $\xi \in \Omega^3(M;\mathfrak{g}_E)$ we define the natural pairing as

$$\xi(X) = 2\int_M \kappa(X,\xi).$$

Then for $X, Y \in \Omega^1(M;\mathfrak{g}_E)$ and $\xi, \eta \in \Omega^3(M;\mathfrak{g}_E)$ we have

$$\langle X+\xi, Y+\eta\rangle = \int_M \kappa(X,\eta) + \kappa(Y,\xi) = \int_M \kappa(X,\eta) - \kappa(\xi,Y) = \int_M \kappa(X+\xi, Y+\eta)_{Ch}$$

where $\kappa(\cdot,\cdot)_{Ch}$ indicates that one uses κ to pair elements in $\mathfrak{g}_E$ and the Chevalley pairing on forms to obtain a top degree form.

We denote by $\widetilde{G}$ the gauge group of E and by $\widetilde{\mathfrak{g}} = \Omega^0(M;\mathfrak{g}_E)$ its Lie algebra. The infinitesimal generator corresponding to $\gamma \in \Omega^0(M;\mathfrak{g}_E)$ at a point $D \in \mathbb{A}$ is just the vector $D\gamma \in \Omega^1(M;\mathfrak{g}_E)$ and we can extend this action to form a lifted action as in [7]:

$$\widetilde{\Psi} : \Omega^0(M;\mathfrak{g}_E) \longrightarrow \mathbb{TA} \qquad \widetilde{\Psi}(\gamma)|_D = D^H\gamma = D\gamma + H\wedge\gamma,$$

as long as there are no infinitesimal symmetries, i.e., as long as $D : \Omega^0(M;\mathfrak{g}_E) \longrightarrow \Omega^1(M;\mathfrak{g}_E)$ has trivial kernel. Therefore, *from this point onwards we only consider connections for which* $D : \Omega^0(M;\mathfrak{g}_E) \longrightarrow \Omega^1(M;\mathfrak{g}_E)$ *has trivial kernel.* If E is a simple $SU(n)$-bundle then that is the case for all ASD connections.

Next we add a moment map to this action. Our (equivariant) moment map takes values on the $\widetilde{G}$-module $\mathfrak{h}^* = \Omega^2_+(M;\mathfrak{g}_E)$, the space of self-dual 2-forms:

$$\mu : \mathbb{A} \longrightarrow \Omega^2_+(M;\mathfrak{g}_E) \qquad \mu(D) = (F_D)_+,$$

where F_D is the curvature of the connection D and $(\cdot)_+$ indicates projection onto the space of self dual forms, $\Omega^\bullet_+(M;\mathfrak{g}_E)$, which, for 2-forms in 4 dimensions, agrees with the usual self dual 2-forms. If we let $\mathfrak{a}$ be the sum $\widetilde{\mathfrak{g}}\oplus\mathfrak{h}$, $\mathfrak{a}$ becomes a Courant algebra if we endow it with the hemisemidirect product:

$$[\![(\gamma_1,\lambda_1),(\gamma_2,\lambda_2)]\!] = ([\gamma_1,\gamma_2],\gamma_1\cdot\lambda_2).$$

Then the maps $\widetilde{\Psi}$ and μ together give rise to an extended action given infinitesimally by map of Courant algebras:

$$\Psi:\mathfrak{a}\longrightarrow\Gamma(\mathbb{TA})\qquad \Psi(\gamma,\lambda)|_D = D^H\gamma + d\langle\mu,\lambda\rangle.$$

We notice that $\Omega^0(M;\mathfrak{g}_E)+\Omega^2_+(M;\mathfrak{g}_E)$ is isomorphic to $\Omega^{ev}_+(M;\mathfrak{g}_E)$ via the map

$$\gamma+\lambda\mapsto\gamma+\lambda+\star\gamma,\qquad \gamma\in\Omega^0(M;\mathfrak{g}_E),\lambda\in\Omega^2_+(M;\mathfrak{g}_E),$$

so we can use $\mathfrak{a}=\Omega^{ev}_+(M;\mathfrak{g}_E)$ and then the extended action is given simply by

$$\Psi:\Omega^{ev}_+(M;\mathfrak{g}_E)\longrightarrow\Omega^{od}(M;\mathfrak{g}_E)\qquad \Psi(\alpha)|_D = D^H\alpha.$$

Following the reduction procedure, the reduced manifold, $\mathcal{M}$, is obtained by taking the quotient of $\mu^{-1}(0)$ by the action of the gauge group. In this case, $\mu^{-1}(0)$ consists of the space of anti self-dual connections and hence $\mathcal{M}$ is simply the moduli space of instantons.

The reduction procedure also produces a specific Courant algebroid over the reduced manifold. Namely, if we let $\mathbb{K}$ be the bundle generated by $\Psi(\mathfrak{a})$ and $\mathbb{K}^\perp$ its orthogonal complement with respect to the natural pairing, then the space of $\widetilde{G}$-invariant sections of $\mathbb{K}^\perp/\mathbb{K}$ over $\mu^{-1}(0)$ inherits a bracket and a nondegenerate pairing which make the quotient bundle $(\mathbb{K}^\perp/\mathbb{K})/\widetilde{G}$ into a Courant algebroid over $\mathcal{M}$. Notice that at a specific anti self-dual connection D, $\mathbb{K}$ is the image of $\Omega^{ev}_+(M;\mathfrak{g}_E)$ by D^H. If we let $D^H_+:\Omega^{od}(M;\mathfrak{g}_E)\longrightarrow\Omega^{ev}_+(M;\mathfrak{g}_E)$ be the composition of D^H with projection onto the SD-forms, then a simple integration by parts shows that $\mathbb{K}^\perp|_D$ is the space of D^H_+-closed forms and hence the reduced Courant algebroid is the degree one cohomology of the following elliptic complex:

$$\{0\}\longrightarrow\Omega^{ev}_+(M;\mathfrak{g}_E)\xrightarrow{D^H}\Omega^{od}(M;\mathfrak{g}_E)\xrightarrow{D^H_+}\Omega^{ev}_+(M;\mathfrak{g}_E)\longrightarrow\{0\}. \tag{6.1}$$

In a way, this is the double of the usual elliptic complex describing the tangent space to M:

$$\{0\}\longrightarrow\Omega^0(M;\mathfrak{g}_E)\xrightarrow{D}\Omega^1(M;\mathfrak{g}_E)\xrightarrow{D_+}\Omega^2_+(M;\mathfrak{g}_E)\longrightarrow\{0\}. \tag{6.2}$$

Earlier we added the assumption that (6.2) above has no cohomology in degree zero. From now on we also add the assumption that this complex has no cohomology in degree two, so that D corresponds to a smooth point in $\mathcal{M}$ and hence the dimension of $\mathcal{M}$ is given by the index of D, which is a topological invariant due to the Atiyah-Singer index theorem. Further, in this case, the cohomology of (6.1) also concentrates in the middle term:

$$H^{od}_{D^H}(M;\mathfrak{g}_E)=\frac{\ker(D^H_+:\Omega^{od}(M;\mathfrak{g}_E)\longrightarrow\Omega^{ev}_+(M;\mathfrak{g}_E)}{\mathrm{Im}\,(D^H:\Omega^{ev}_+(M;\mathfrak{g}_E)\longrightarrow\Omega^{od}(M;\mathfrak{g}_E)}.$$

Since M has an SKT structure, we can also endow $\mathbb{A}$ with an SKT structure. Firstly we let $\star$ be the generalized metric: Indeed, in four dimensions $\star^2 = \mathrm{Id}$, spin invariance of the Chevalley pairing means that $\star$ is orthogonal (in even dimensions) and by design we have (c.f. (5.1))

$$\langle \varphi, \star\varphi \rangle = \int_M \kappa(\varphi, \star\varphi)_{Ch} > 0, \qquad \text{for } \varphi \in \Omega^{od}(M; \mathfrak{g}_E)\backslash\{0\},$$

so $\star$ is a generalized metric and for this metric V_+ is the space of SD odd forms, $\Omega^{od}_+(M; \mathfrak{g}_E) = (\mathcal{W}^2 + \mathcal{W}^{-2}) \cap \Omega^{od}(M; \mathfrak{g}_E)$, and V_- is the space of ASD odd forms, $\Omega^{od}_-(M; \mathfrak{g}_E) = \mathcal{W}^0 \cap \Omega^{od}(\mathcal{M}; \mathfrak{g}_E)$.

For complex structure, we let $\mathbb{I} = e^{\frac{\pi\mathcal{I}}{2}}$ as in Lemma 1.10. Then that same lemma implies that $\mathbb{I}^2|_{V_+} = -\star = -\mathrm{Id}$. Since this structure is independent of the point $D \in \mathbb{A}$, it is constant and hence integrable. The spaces $V_+^{1,0}$ and $V_+^{0,1}$ can be easily described: since $\mathcal{I} = \mathbb{I}$ on $W^{\pm 2}$ we have $V_+^{1,0} = \mathcal{W}^2 \cap \Omega^{od}(M; \mathfrak{g}_E)$ and $V_+^{0,1} = \mathcal{W}^{-2} \cap \Omega^{od}(M; \mathfrak{g}_E)$.

Further, since both $\star$ and $\mathbb{I}$ are constant and the gauge group acts by translations on each of its orbits, this SKT structure is invariant by the action of the gauge group. Now, we are in position to use the SKT reduction theorem:

Theorem 6.1 (SKT Reduction Theorem [7])**.** *Let $\Psi : \mathfrak{a} \longrightarrow \Gamma(\mathbb{T}\mathbb{A})$ be an extended action with moment map μ preserving an SKT structure $(\mathcal{G}, \mathcal{I})$ on $\mathbb{A}$ as above and let $P = \mu^{-1}(0)$. If the underlying group action on P is free and proper and 0 is a regular value of the moment map, then $\mathcal{M}_{red} = P/G$ is smooth and the SKT structure on $\mathbb{A}$ reduces to an SKT structure on $\mathcal{M}_{red}$ if and only if $\mathbb{K}^\perp \cap V_+|_P$ is invariant under $\mathcal{I}$.*

As we observed earlier, at a point $D \in \mathbb{A}$, $\mathbb{K}^\perp$ corresponds to the D^H_+-closed odd forms and hence $\mathbb{K}^\perp \cap V_+$ corresponds to the D^H_+-closed SD odd forms, that is the SD, D^H-harmonic odd forms and hence according to the theorem to prove that the moduli space of instantons inherits an SKT from M, we must prove that the space of SD, D^H-harmonic odd forms is invariant under $\mathbb{I}$. To achieve this, one must develop Hodge theory for forms with coefficients. In this case, the condition that the connection is anti self-dual, allows us to achieve the result.

Indeed, similarly to the flat case, we can split D^H as a sum of three operators:

$$D^H = \delta_+^N + \overline{\delta_+^N} + \not{D}_-^H,$$

$$\delta_+^N : \mathcal{W}^k \longrightarrow \mathcal{W}^{k+2} \qquad \overline{\delta_+^N} : \mathcal{W}^k \longrightarrow \mathcal{W}^{k-2} \qquad \not{D}_-^H : \mathcal{W}^k \longrightarrow \mathcal{W}^k.$$

Indeed, locally $D^H = d^H + A$, for some $A \in \Omega^1(M; \mathfrak{g}_E)$ and hence the splitting of d^H together with the splitting of A into its $V_+^{1,0}$, $V_+^{0,1}$ and V_- components gives the desired decomposition of D^H. Just as in Section 5.2, integration by parts gives that $(\delta_+^N)^* = -\overline{\delta_+^N}$ and $\not{D}_-^{H*} = \not{D}_-^H$.

Now, let $\varphi \in (\mathcal{W}^2 \oplus \mathcal{W}^{-2}) \cap \Omega^{od}(M; \mathfrak{g}_E)$. Then the D^H-Laplacian computed on φ is given by

$$\begin{aligned} \triangle_H \varphi &= (D^{H*} D_+^H + D^H D_+^{H*})\varphi \\ &= (-\delta_+^N - \overline{\delta_+^N} + \not{D}_-^H)\not{D}_-^H + (\delta_+^N + \overline{\delta_+^N} + \not{D}_-^H)\not{D}_-^H \varphi \\ &= 2(\not{D}_-^H)^2 \varphi. \end{aligned}$$

Hence the Laplacian leaves the spaces $\mathcal{W}^2 \cap \Omega^{od}$ and $\mathcal{W}^{-2} \cap \Omega^{od}$ invariant. Therefore we can decompose the space of harmonic SD odd forms into two spaces $\mathcal{H}^{\pm 2} = \ker(\triangle_H) \cap \mathcal{W}^{\pm 2}$. $\mathbb{I}$ acts as multiplication by i on $\mathcal{W}^2$ and by $-i$ on $\mathcal{W}^{-2}$, so either way it preserves the intersection $\ker(\triangle_H) \cap \mathcal{W}^{\pm 2}$ and hence it preserves the space of SD harmonic odd forms. According to Theorem 6.1, this means that the SKT structure from $\mathbb{A}$ reduces to an SKT structure on $\mathcal{M}$ so we have re-obtained the following result, originally due to Lübke and Teleman [28]:

Theorem 6.2. *Let $(M, I, [g])$ be a compact conformal Hermitian 4-manifold, let E be a bundle over M whose structure group is compact with Lie algebra $\mathfrak{g}$ and let $\mathfrak{g}_E$ be the adjoint bundle over M. Let $\mathcal{M}_s$ be the quotient of the space*

$$\widetilde{P} = \{D \in \mathbb{A} : (F_D)_+ = 0;\ H^0_D(M; \mathfrak{g}_E) = H^2_D(M; \mathfrak{g}_E) = \{0\}\},$$

by the action of the gauge group, i.e., $\mathcal{M}_s$ is the smooth locus of the moduli space of instantons on E. Then $\mathcal{M}_s$ has an SKT structure induced by the unique SKT structure on M in the conformal class $[g]$.

7 Lie algebroid differentials and other operators

Lie algebroids occur naturally in generalized geometry. Notably, if $\mathcal{J}$ is a generalized complex structure and L its $+i$-eigenspace, then L is a Lie algebroid and the corresponding degree two Lie algebroid cohomology, $H^2(M; L^*)$, governs the infinitesimal deformations of $\mathcal{J}$. As we will see next section, similar claims also hold in the SKT case, so it is particularly relevant to develop the theory of Lie algebroids related to SKT structures. However, since some of the generalized complex structures we consider are not integrable, some bundles we encounter are not Lie algebroids. Yet some of those "not Lie algebroids" are important in the study of deformations, so we must develop the relevant part of the theory of these spaces as well.

7.1 Lie algebroids

In this section we introduce two natural Lie algebroids: one associated to an SKT structure and one to a generalized complex extension of an SKT structure. Then we study the relation between the differential operators defined in the previous section and the natural operators defined on these Lie algebroids.

If (M, H) has a positive SKT structure $(\mathcal{G}, \mathcal{I})$, by definition $V_+^{1,0}$ is involutive, hence it is a Lie algebroid over M. Further $V_+^{0,1}$ can be identified with the dual of $V_+^{1,0}$ using the natural pairing, which is nondegenerate on V_+, hence we get a corresponding differential

$$\overline{\partial}_+ : \Gamma(\wedge^k V_+^{0,1}) \longrightarrow \Gamma(\wedge^{k+1} V_+^{0,1}). \tag{7.1}$$

Similarly, given a generalized complex extension $\mathcal{J}_1$ of an SKT structure, L_1, the $+i$-eigenspace of $\mathcal{J}_1$, is also a Lie algebroid and we get a differential on the exterior algebra of the dual $d_{L_1} : \Gamma(\wedge^j \overline{L_1}) \longrightarrow \Gamma(\wedge^{j+1} \overline{L_1})$

Since $\mathcal{J}_2 = \mathcal{G}\mathcal{J}_1$ commutes with $\mathcal{J}_1$, it preserves the eigenspaces of $\mathcal{J}_1$ and hence gives rise to a complex structure on L_1. In fact, it splits L_1 in two subspaces of complex dimension n: $V_+^{1,0} = L_1 \cap V_+$ and $V_-^{1,0} = L_1 \cap V_-$, hence $\wedge^\bullet \overline{L_1}$ gets a bigrading:

$$\wedge^{k,l}\overline{L_1} = \wedge^k V_+^{0,1} \otimes \wedge^l V_-^{0,1}.$$

Proposition 7.1. *Let $\mathcal{J}_1$ be a generalized complex extension of a positive SKT structure. Then*

$$d_{L_1} : \Gamma(\wedge^{k,l}\overline{L_1}) \longrightarrow \Gamma(\wedge^{k+1,l}\overline{L_1} \oplus \wedge^{k,l+1}\overline{L_1} \oplus \wedge^{k-1,l+2}\overline{L_1})$$

Further,

- *The component*

$$\mathcal{N} = \pi_{k-1,l+2} \circ d_{L_1} : \Gamma(\wedge^{k,l}\overline{L_1}) \longrightarrow \Gamma(\wedge^{k-1,l+2}\overline{L_1})$$

corresponds to the Clifford action of the Nijenhuis tensor $N \in \Gamma(\wedge^2 V_-^{0,1} \otimes V_+^{1,0})$ introduced in (3.2);

- *The component*

$$\pi_{k+1,0} \circ d_{L_1} : \Gamma(\wedge^{k,0}\overline{L_1}) \longrightarrow \Gamma(\wedge^{k+1,0}\overline{L_1})$$

corresponds to the operator $\overline{\partial}_+$ introduced in (7.1).

Proof. Since $V_+^{1,0}$ is involutive and $\wedge^{0,1}\overline{L_1} = V_-^{0,1}$ is the space of forms annihilating $V_+^{1,0}$ in $\overline{L_1}$, we have

$$d_{L_1} : \Gamma(\wedge^{0,1}\overline{L_1}) \longrightarrow \Gamma(\wedge^{0,2}\overline{L_1} \oplus \wedge^{1,1}\overline{L_1}).$$

On the other hand, there are no hypothesis about integrability of $V_-^{1,0}$, hence

$$d_{L_1} : \Gamma(\wedge^{1,0}\overline{L_1}) \longrightarrow \Gamma(\wedge^{0,2}\overline{L_1} \oplus \wedge^{1,1}\overline{L_1} \oplus \wedge^{2,0}\overline{L_1}).$$

Using the Leibniz rule, we have

$$d_{L_1} : \Gamma(\wedge^{k,l}\overline{L_1}) \longrightarrow \Gamma(\wedge^{k+1,l}\overline{L_1} \oplus \wedge^{k,l+1}\overline{L_1} \oplus \wedge^{k-1,l+2}\overline{L_1}).$$

One can easily check that $\mathcal{N}$ is just the Nijenhuis tensor corresponding to the complex structure $\mathcal{J}_2|_{L_1}$, i.e.,

$$\mathcal{N} \in \wedge^2 V_-^{0,1} \otimes V_+^{1,0}, \qquad \mathcal{N}(X,Y) = -[\![X,Y]\!]_+, \qquad \text{for all } X, Y \in \Gamma(V_-^{1,0}),$$

where $\cdot_+$ indicates projection onto V_+ and this is precisely how the operator N introduced in (3.2) is defined, hence these two operators agree.

Finally, for $\alpha \in \Gamma(\wedge^{k,0}\overline{L_1}) = \Gamma(\wedge^k V_+^{0,1})$, $\pi_{k+1,0} \circ d_{L_1}\alpha$ is determined by its values on $k+1$ elements $v_0, \cdots, v_k \in V_+^{1,0}$:

$$\begin{aligned}(\pi_{k+1,0} \circ d_{L_1}\alpha)(v_0, \cdots, v_k) &= (d_{L_1}\alpha)(v_0, \cdots, v_k)\\ &= \sum_i (-1)^i \mathcal{L}_{\pi_T v_i} \alpha(v_0, \cdots, \hat{v}_i, \cdots, v_k) +\\ &+ \sum_{i,j} (-1)^{i+j} \alpha([\![v_i, v_j]\!], v_0, \cdots, \hat{v}_i \cdots, \hat{v}_j, \cdots, v_k)\\ &= (\overline{\partial}_+\alpha)(v_0, \cdots, v_k),\end{aligned}$$

where we have simply used the definitions of the operators d_{L_1} and $\overline{\partial}_+$. □

This allows us to define operators

$$\begin{aligned}\overline{\partial}_+ &: \Gamma(\wedge^{k,l}\overline{L_1}) \longrightarrow \Gamma(\wedge^{k+1,l}\overline{L_1});\\ \overline{\partial}_- &: \Gamma(\wedge^{k,l}\overline{L_1}) \longrightarrow \Gamma(\wedge^{k,l+1}\overline{L_1});\end{aligned}$$

and the condition $d_{L_1}^2 = 0$ gives:

Corollary 7.2. Let $\overline{\partial}_+$, $\overline{\partial}_-$ and $\mathcal{N}$ be the operators defined above. Then

$$\overline{\partial}_+^2 = \mathcal{N}^2 = 0 \qquad \text{and} \qquad \overline{\partial}_-^2 = -\{\overline{\partial}_+, \mathcal{N}\}.$$

Finally, the operator d_{L_1} is related to the operator $\overline{\partial}_{\mathcal{J}_1}$ via (see [21, 27]),

$$\overline{\partial}_{\mathcal{J}_1}(\alpha \cdot \varphi) = (d_{L_1}\alpha) \cdot \varphi + (-1)^{|\alpha|}\alpha \cdot \overline{\partial}_{\mathcal{J}_1}\varphi.$$

So, if we take $\alpha \in \Gamma(\wedge^{k,l}\overline{L_1})$ and $\varphi \in \mathcal{U}^{p,q}$ we can split the equation above in three components which give the following relations:

Proposition 7.3. *For* $\alpha \in \Gamma(\wedge^{\bullet}\overline{L_1})$ *and* $\varphi \in \Omega^{\bullet}(M;\mathbb{C})$ *we have*

$$\begin{aligned}\{\overline{\delta}_+, \alpha\} \cdot \varphi &= (\overline{\partial}_+\alpha) \cdot \varphi;\\ \{\overline{\delta}_-, \alpha\} \cdot \varphi &= (\overline{\partial}_-\alpha) \cdot \varphi;\\ \{N, \alpha\} \cdot \varphi &= \mathcal{N}(\alpha) \cdot \varphi.\end{aligned}$$

7.2 Further operators

Proposition 7.3 relates the Lie algebroid operators to the graded commutator of the action of an element $\alpha \in \Gamma(\wedge^{\bullet}\overline{L_1})$ and the operators $\overline{\delta}_+$, $\overline{\delta}_-$ and N. In an SKT manifold, one can define more operators that can be obtained this way but which do not come from Lie algebroids. The reason to consider such operators will be made clear next section when studying the deformation and stability problem. There we will see that the correct bundle to consider is L_2, the $+i$-eigenspace of $\mathcal{J}_2$. Since L_2 is not a Lie algebroid we are forced to consider differential operators associated to brackets which do not satisfy Jacobi, and hence do not square to zero, but we don't let this mishap dampen our sentiment.

As before, we let $\mathcal{J}_1$ be a generalized complex extension of the SKT structure and inspired by Proposition 7.3 we define

$$\begin{aligned}\overline{\partial}_\pm &: \mathrm{Clif}(\mathbb{T}_{\mathbb{C}}M) \longrightarrow \mathrm{Diff}(\Omega^{\bullet}(M;\mathbb{C})) & \overline{\partial}_\pm\alpha &= \{\overline{\delta}_\pm, \alpha\};\\ \partial_\pm &: \mathrm{Clif}(\mathbb{T}_{\mathbb{C}}M) \longrightarrow \mathrm{Diff}(\Omega^{\bullet}(M;\mathbb{C})) & \partial_\pm\alpha &= \{\delta_\pm, \alpha\};\\ \overline{\mathcal{N}} &: \mathrm{Clif}(\mathbb{T}_{\mathbb{C}}M) \longrightarrow \mathrm{Clif}(\mathbb{T}_{\mathbb{C}}M) & \overline{\mathcal{N}}\alpha &= \{\overline{N}, \alpha\};\end{aligned} \tag{7.2}$$

where $\mathrm{Diff}(\Omega^{\bullet}(M;\mathbb{C}))$ is the vector space of differential operators on forms which is itself a $\mathbb{Z}_2$-graded Lie algebra with graded commutator as the bracket.

We denote $\Gamma(\wedge^{k,l}\overline{L_2}) = \Gamma(\wedge^k V_+^{0,1} \otimes \wedge^l V_-^{1,0})$. Next we summarize the main properties of these operators.

Proposition 7.4. *The restriction of $\overline{\mathcal{N}}$ to $\wedge^\bullet\overline{L_2}$ vanishes and*

$$\overline{\partial}_+ : \Gamma(\wedge^k V_+^{0,1} \otimes \wedge^l V_-^{1,0} \otimes \wedge^m V_-^{0,1}) \longrightarrow \Gamma(\wedge^{k+1} V_+^{0,1} \otimes \wedge^l V_-^{1,0} \otimes \wedge^m V_-^{0,1}); \tag{7.3}$$

$$\partial_- : \Gamma(\wedge^{k,l}\overline{L_2}) \longrightarrow \Gamma(\wedge^{k,l+1}\overline{L_2}). \tag{7.4}$$

Further, the following relations and their complex conjugates hold:

$$\overline{\partial}_+^2 = \overline{\mathcal{N}}^2 = 0; \qquad \{\overline{\partial}_+, \overline{\mathcal{N}}\} = 0;$$

$$\partial_-^2 = -\{\overline{\mathcal{N}}, \overline{\partial}_+\}; \qquad \{\overline{\mathcal{N}}, \partial_-\} = 0; \qquad \{\overline{\partial}_-, \overline{\partial}_+\} = 0; \qquad \{\overline{\partial}_+, \partial_-\} = -\{\overline{\partial}_-, \overline{\mathcal{N}}\};$$

$$\{\partial_+, \overline{\partial}_+\} + \{\partial_-, \overline{\partial}_-\} + \{\mathcal{N}, \overline{\mathcal{N}}\} = 0.$$

Proof. The relations between the operators follow from the corresponding relations for $\overline{\delta}_+$, δ_- and $\overline{N}$ and the fact that the vector space of operators on forms are a $\mathbb{Z}_2$-graded Lie algebra with Lie bracket given by the graded commutator of operators. If we let $\alpha \in \mathrm{Clif}(\mathbb{T}_\mathbb{C}M)$, using the graded Jacobi identity we have

$$\{d^H, \{d^H, \alpha\}\} = \{\{d^H, d^H\}, \alpha\} - \{d^H, \{d^H, \alpha\}\} = -\{d^H, \{d^H, \alpha\}\} = 0,$$

where we have used that $\{d^H, d^H\} = 2(d^H)^2 = 0$. Hence, splitting d^H into its six components, taking $\alpha \in \Gamma(\wedge^j V_+^{1,0} \otimes \wedge^k V_+^{0,1} \otimes \wedge^l V_-^{1,0} \otimes \wedge^m V_-^{0,1})$, applying this operator to elements in $\mathcal{U}^{p,q}$ and imposing the vanishing of the different components of the result we get the relations stated in the theorem. For example the vanishing of the component in $\mathcal{U}^{p+(j-k)+(l-m)+2, q+(j-k)-(l-m)}$ gives

$$\{\delta_-, \{\overline{\delta}_+, \alpha\}\} + \{\overline{\delta}_+, \{\delta_-, \alpha\}\} + \{\overline{\delta}_-, \{\overline{N}, \alpha\}\} + \{\overline{N}, \{\overline{\delta}_-, \alpha\}\} = 0.$$

That is

$$\partial_-\overline{\partial}_+ + \overline{\partial}_+\partial_- + \overline{\partial}_-\overline{\mathcal{N}} + \overline{\mathcal{N}}\partial_- = 0.$$

Which is the same as

$$\{\overline{\partial}_+, \partial_-\} + \{\overline{\partial}_-, \overline{\mathcal{N}}\} = 0$$

Since $\overline{N} \in \Gamma(\wedge^3\overline{L_2})$ and $\overline{L_2}$ is isotropic, $\{\overline{N}, \alpha\}$ vanishes for every $\alpha \in \Gamma(\wedge^\bullet\overline{L_2})$.

Next we will prove that (7.3) and (7.4) hold. Firstly, we prove that when restricted to the domains indicated in (7.3) and (7.4) $\overline{\partial}_+$ and ∂_- actually take values in the Clifford algebra of $\mathbb{T}M$. Indeed, for $\alpha \in \mathrm{Clif}(\mathbb{T}M)$, $\varphi \in \mathcal{U}^{p,q}$ and $f \in C^\infty(M)$ we decompose df into its $V_+^{1,0}$, $V_+^{0,1}$, $V_-^{1,0}$ and $V_-^{0,1}$ components, say, $\partial_+ f$, $\overline{\partial}_+ f$, $\partial_- f$ and $\overline{\partial}_- f$, respectively and compute

$$\begin{aligned}
\{d^H, \alpha\}(f\varphi) &= d^H \alpha f\varphi - (-1)^{|\alpha|}\alpha d^H(f\varphi) \\
&= df \wedge \alpha\varphi + f d^H\alpha \cdot \varphi - (-1)^{|\alpha|}\alpha(df \wedge \varphi + f d^H\varphi) \\
&= (\partial_- f + \overline{\partial}_- f + \partial_+ f + \overline{\partial}_+ f) \cdot \alpha\varphi + f d^H \alpha\varphi \\
&- (-1)^{|\alpha|}\alpha((\partial_- f + \overline{\partial}_- f + \partial_+ f + \overline{\partial}_+ f) \cdot \varphi + f d^H\varphi). \\
&= f\{d^H\alpha\}\varphi + (\partial_- f + \overline{\partial}_- f + \partial_+ f + \overline{\partial}_+ f) \cdot \alpha\varphi \\
&- (-1)^{|\alpha|}\alpha((\partial_- f + \overline{\partial}_- f + \partial_+ f + \overline{\partial}_+ f) \cdot \varphi).
\end{aligned} \tag{7.5}$$

If $\alpha \in \Gamma(\wedge^k V_+^{0,1} \otimes \wedge^l V_-^{1,0} \otimes \wedge^m V_-^{0,1})$, then $\overline{\partial}_+ f$ graded commutes with α and hence if we compose $\{d^H, \alpha\}$ with the projection onto $\mathcal{U}^{p-k+l-m-1,q-k-l+m-1}$ we have

$$\pi_{\mathcal{U}^{p-k+l-m-1,q-k-l+m-1}} \circ \{d^H, \alpha\} : \mathcal{U}^{p,q} \longrightarrow \mathcal{U}^{p-k+l-m-1,q-k-l+m-1}$$

is C^∞-linear, that is, it is a tensor. Further, using the decomposition d^H into its six components, as in Theorem 3.6, we see that the tensor above corresponds to:

$$\overline{\partial}_+ \alpha = \{\overline{\delta}_+, \alpha\} : \mathcal{U}^{p,q} \longrightarrow \mathcal{U}^{p-k+l-m-1,q-k-l+m-1}$$

Since the action of $\mathrm{Clif}(\mathbb{T}M)$ establishes an isomorphism between $\mathrm{Clif}(\mathbb{T}M)$ and the space of endomorphisms of $\wedge^\bullet T^*M$, the tensor $\overline{\partial}_+\alpha$ corresponds to an element in $\mathrm{Clif}(\mathbb{T}M)$:

$$\overline{\partial}_+ : \Gamma(\wedge^k V_+^{0,1} \otimes \wedge^l V_-^{1,0} \otimes \wedge^m V_-^{0,1}) \longrightarrow \mathrm{Clif}(\mathbb{T}_{\mathbb{C}}M)$$

Similarly, if we require α to lie in $\Gamma(\wedge^{k,l}\overline{L_2})$ the same argument gives us the result for ∂_-. Indeed, in this setting, $\partial_- f$ also graded commutes with α in (7.5), hence we get another tensor, this time by composing $\{d^H, \alpha\}$ with the projection onto $\mathcal{U}^{p-k+l-m+1,q-k-l+m-1}$, that is, we get

$$\partial_- : \Gamma(\wedge^{k,l}\overline{L}) \longrightarrow \mathrm{Clif}(\mathbb{T}_{\mathbb{C}}M). \tag{7.6}$$

Now we deal with $\overline{\partial}_+$.

Lemma 7.5. Let $(M, \mathcal{G}, \mathcal{I})$ be a positive SKT manifold and $\mathcal{J}$ be a generalized complex extension of the SKT structure. The vector bundle $V_-^{1,0}$ is a representation of the Lie algebroid $V_+^{1,0}$ if endowed with the connection

$$\nabla : \Gamma(V_+^{1,0} \otimes V_-^{1,0}) \longrightarrow \Gamma(V_-^{1,0}); \qquad \nabla_v w = \pi_{V_-^{1,0}}([\![v, w]\!]),$$

where $v \in \Gamma(V_+^{1,0})$ and $w \in \Gamma(V_-^{1,0})$. That is, for $v, v_1, v_2 \in \Gamma(V_+^{1,0})$, $w \in \Gamma(V_-^{1,0})$ and $f \in C^\infty(M)$ we have

$$\nabla_{fv} w = f\nabla_v w, \qquad \nabla_v fw = (\mathcal{L}_{\pi_T(v)} f)w + f\nabla_v w$$

$$\nabla_{v_1}\nabla_{v_2} - \nabla_{v_2}\nabla_{v_1} - \nabla_{[\![v_1,v_2]\!]} = 0.$$

Proof. The first two identities follow from basic properties of the Courant bracket:

$$\nabla_{fv} w = \pi_{V_-^{1,0}}([\![fv, w]\!]) = \pi_{V_-^{1,0}}(f[\![v, w]\!] - (\mathcal{L}_{\pi_T(w)} f)v) = \pi_{V_-^{1,0}}(f[\![v, w]\!]) = f\nabla_v w.$$

$$\nabla_v fw = \pi_{V_-^{1,0}}([\![v, fw]\!]) = \pi_{V_-^{1,0}}(f[\![v, w]\!] + (\mathcal{L}_{\pi_T(v)} f)w) = f\nabla_v w + (\mathcal{L}_{\pi_T(v)} f)w.$$

Now we prove flatness. Let us first consider the term $\nabla_{v_1}\nabla_{v_2} w$. We decompose $[\![v_2, w]\!]$ into two components, according to the decomposition $L_1 = V_+^{1,0} \oplus V_-^{1,0}$ and $\nabla_{v_2} w$ is the component in $V_-^{1,0}$:

$$[\![v_2, w]\!] = u_+^{1,0} + \nabla_{v_2} w, \qquad u_+^{1,0} \in \Gamma(V_+^{1,0}).$$

Since $V_+^{1,0}$ is involutive, we see that

$$[\![v_1, u_+^{1,0}]\!] \in V_+^{1,0}.$$

Hence the $V_-^{1,0}$ component of $[\![v_1, \nabla_{v_2} w]\!]$ is the $V_-^{1,0}$ component of $[\![v_1, [\![v_2, w]\!]]\!]$, that is

$$\nabla_{v_1}\nabla_{v_2} w = \pi_{V_-^{1,0}}([\![v_1, [\![v_2, w]\!]]\!]).$$

Therefore we have

$$\nabla_{v_1}\nabla_{v_2} w - \nabla_{v_2}\nabla_{v_1} w - \nabla_{[\![v_1,v_2]\!]} w = \pi_{V_-^{1,0}}([\![v_1, [\![v_2, w]\!]]\!] - [\![v_2, [\![v_1, w]\!]]\!] - [\![[\![v_1, v_2]\!], w]\!]) = 0,$$

where in the last equality we used the fact that the Courant bracket satisfies the Jacobi identity. □

Since ∇ is a connection on $V_-^{1,0}$, we can extend it to the exterior algebra of $V_-^{1,0}$:

$$\nabla : \Gamma(\wedge^k V_-^{1,0}) \longrightarrow \Gamma(V_+^{0,1} \otimes \wedge^k V_-^{1,0});$$

$$\nabla_v \alpha = \{v, \nabla\alpha\} \qquad \text{for all } v \in \Gamma(V_+^{1,0}).$$

Lemma 7.6. For $\alpha \in \Gamma(\wedge^k V_-^{1,0})$ we have

$$\overline{\partial}_+ \alpha = \nabla\alpha.$$

Proof. Indeed, we check that when paired (i.e., graded commuted) with an element $v \in \Gamma(V_+^{1,0})$ these two operators give the same result. Unwinding the definition of $\overline{\partial}_+$, we have that for $\varphi \in \mathcal{U}^{p,q}$ $\{v, \overline{\partial}_+\alpha\}\varphi$ is the $\mathcal{U}^{p+k,q-k}$ component of $\{v, \{d^H, \alpha\}\}\varphi$. For α decomposable, $\alpha = \alpha_1 \wedge \cdots \wedge \alpha_k$ we have

$$\begin{aligned}
\{v, \overline{\partial}_+\alpha\}\varphi &= \pi_{\mathcal{U}^{p+k,q-k}}(\{v, \{d^H, \alpha_1 \wedge \cdots \wedge \alpha_k\}\}\varphi) \\
&= \pi_{\mathcal{U}^{p+k,q-k}}(\sum_{i=1}^k \alpha_1 \wedge \cdots \wedge [\![v, \alpha_i]\!] \wedge \cdots \wedge \alpha_k \varphi) \\
&= \sum_{i=1}^k \alpha_1 \wedge \cdots \wedge \pi_{V_-^{1,0}}[\![v, \alpha_i]\!] \wedge \cdots \wedge \alpha_k \varphi \\
&= \sum_{i=1}^k \alpha_1 \wedge \cdots \wedge \{v, \nabla\alpha_i\} \wedge \cdots \wedge \alpha_k \varphi \\
&= \{v, \nabla\alpha\}\varphi,
\end{aligned}$$

where in the second equality we used that the Courant bracket is the derived bracket associated to d^H hence the commutator becomes a Schouten–Nijenhuis type bracket between the sections involved.

Since $\overline{\partial}_+\alpha$ and $\nabla\alpha$ are both elements in $\mathrm{Clif}(\mathbb{T}_{\mathbb{C}}M)$ we have just established that their difference lies in the annihilator of $V_+^{1,0}$ under graded commutator, that is

$$\nabla\alpha - \overline{\partial}_+\alpha \in \Gamma(\wedge^\bullet V_+^{1,0} \otimes \wedge^\bullet V_-).$$

And we have that when acting in $\mathcal{U}^{p,q}$

$$\nabla\alpha - \overline{\partial}_+\alpha : \mathcal{U}^{p,q} \longrightarrow \mathcal{U}^{p+k-1,q-k-1} \subset \mathcal{W}^{p+q-2}$$

as individually both $\overline{\partial}_+\alpha$ and $\nabla\alpha$ have this property. On the other hand, elements in $\wedge^l V_+^{1,0} \otimes \wedge^m V_-$ map $\mathcal{U}^{p,q} \subset \mathcal{W}^{p+q}$ into $\mathcal{W}^{p+q+l}$, so the difference $\nabla\alpha - \overline{\partial}_+\alpha$ vanishes and the result follows. □

Proof of (7.3). Now, let $\alpha^{k,l} \in \Gamma(\wedge^k V_+^{0,1} \otimes \wedge^l V_-^{0,1}) = \Gamma(\wedge^{k,l}\overline{L_1})$ and $\alpha_-^{m,0} \in \Gamma(\wedge^m V_-^{1,0})$. Then for a form φ we have

$$\begin{aligned}
\overline{\partial}_+(\alpha^{k,l}\alpha_-^{m,0}) \cdot \varphi &= \{\overline{\delta}_+, \alpha^{k,l}\alpha_-^{m,0}\}\varphi \\
&= \overline{\delta}_+(\alpha^{k,l}\alpha_-^{m,0}\varphi) + (-1)^{k+l+m}\alpha^{k,l}\alpha_-^{m,0}\overline{\delta}_+\varphi \\
&= (\overline{\partial}_+\alpha^{k,l})\alpha_-^{m,0}\varphi + (-1)^{k+l}\alpha^{k,l}\overline{\delta}_+(\alpha_-^{m,0}\varphi) + (-1)^{k+l+m}\alpha^{k,l}\alpha_-^{m,0}\overline{\delta}_+\varphi \\
&= (\overline{\partial}_+\alpha^{k,l})\alpha_-^{m,0}\varphi + (-1)^{k+l}\alpha^{k,l}(\overline{\partial}_+\alpha_-^{m,0})\varphi \\
&= ((\overline{\partial}_+\alpha^{k,l})\alpha_-^{m,0} + (-1)^{k+l}\alpha^{k,l}(\overline{\partial}_+\alpha_-^{m,0}))\varphi.
\end{aligned}$$

According to Proposition 7.3, $(\overline{\partial}_+\alpha^{k,l})\alpha_-^{m,0} \in \Gamma(\wedge^{k+1}V_+^{0,1}\otimes\wedge^l V_-^{0,1}\otimes\wedge^m V_-^{1,0})$ and, according to Lemma 7.6, $\alpha^{k,l}(\overline{\partial}_+\alpha_-^{m,0})$ also belongs to this space, hence we have proved (7.3). ☐

Proof of (7.4). Since $\overline{L_2}$ is maximal isotropic, $\wedge^\bullet\overline{L_2} \subset \mathrm{Clif}(\mathbb{T}_\mathbb{C}M)$ can be characterized as

$$\wedge^\bullet\overline{L_2} = \{\beta \in \mathrm{Clif}(\mathbb{T}_\mathbb{C}M) : \{\beta, v\} = 0 \text{ for all } v \in \overline{L_2}\}.$$

Now, given $\alpha \in \Gamma(\wedge^{k,l}\overline{L_2})$, we already know that

$$\partial_-\alpha : \mathcal{U}^{p,q} \longrightarrow \mathcal{U}^{p-k+l+1,q-k-l-1} \tag{7.7}$$

so to prove that $\partial_-\alpha \in \Gamma(\wedge^{k,l+1}\overline{L_2})$ it is enough to prove that $\partial_-\alpha \in \Gamma(\wedge^\bullet\overline{L_2})$ as the only elements in $\Gamma(\wedge^\bullet\overline{L_2})$ with property (7.7) are those in $\Gamma(\wedge^{k,l+1}\overline{L_2})$. So, for $\alpha \in \Gamma(\wedge^{k,l}\overline{L_2})$, $v \in \Gamma(\overline{L_2})$ and $\varphi \in \mathcal{U}^{p,q}$ we compute

$$\begin{aligned}
\{\partial_-\alpha, v\}\varphi &= \{\{\delta_-, \alpha\}, v\}\varphi \\
&= \pi_{\mathcal{U}^{p-k+l,q-k-l-2}}(\{\{d^H, \alpha\}, v\}\varphi) \\
&= \pm\pi_{\mathcal{U}^{p-k+l,q-k-l-2}}([\![\alpha, v]\!]_{SN}\varphi) \\
&= 0,
\end{aligned}$$

where in the third equality we used that $\overline{L_2}$ is isotropic, and hence the derived bracket is just the extension of the Courant bracket to a Schouten–Nijenhuis-like bracket and in the last equality we have used that we have used that the Schouten–Nijenhuis bracket has degree -1, hence it maps $(v, \alpha) \in \overline{L_2} \times \wedge^{k,l}\overline{L_2}$ into $\mathrm{Clif}^{k+l}(\mathbb{T}_\mathbb{C}M)$ and hence it can not map $\mathcal{U}^{p,q}$ to $\mathcal{U}^{p-k+l,q-k-l-2}$ as elements in $\mathrm{Clif}^j(\mathbb{T}_\mathbb{C}M)$ can only change the first and the second index of the bigrading of forms (the 'p' and the 'q') by at most $\pm j$.

☐

As a consequence of Proposition 7.4, we see that although L_2 is not a Lie algebroid, we can still define some cohomology there using $\overline{\partial}_+$.

Definition 7.7. Let $(\mathcal{G}, \mathcal{I})$ be a positive SKT structure on a manifold with 3-form (M, H) and $\mathcal{J}$ be a generalized complex extension of $\mathcal{I}$. The *$\overline{\partial}_+$-cohomology in $\overline{L_2}$*, $H^{\bullet,\bullet}_{\overline{\partial}_+}(M; \overline{L_2})$, is the quotient

$$H^{k,l}_{\overline{\partial}_+}(M; \overline{L_2}) = \frac{\ker(\overline{\partial}_+ : \Gamma(\wedge^{k,l}\overline{L_2}) \longrightarrow \Gamma(\wedge^{k+1,l}\overline{L_2}))}{\mathrm{Im}\,(\overline{\partial}_+ : \Gamma(\wedge^{k-1,l}\overline{L_2}) \longrightarrow \Gamma(\wedge^{k,l}\overline{L_2}))}$$

Finally we observe that all the operators we have studied in this section where obtained with the help of a generalized complex extension of the SKT structure, so, they are not intrinsic to the SKT structure. Yet, one could follow the same road and define

$$\overline{\partial_+^{\mathcal{N}}} : \mathrm{Clif}(\mathbb{T}_{\mathbb{C}}M) \longrightarrow \mathrm{Diff}(\Omega^\bullet(M;\mathbb{C})) \qquad \overline{\partial_+^{\mathcal{N}}}\alpha = \{\overline{\delta_+^N}, \alpha\};$$

$$d_- : \mathrm{Clif}(\mathbb{T}_{\mathbb{C}}M) \longrightarrow \mathrm{Diff}(\Omega^\bullet(M;\mathbb{C})); \qquad d_-\alpha = \{\not{D}_-, \alpha\}.$$

By choosing a generalized complex extension to $\mathcal{I}$, the decomposition $\delta_+^N = \delta_+ + N$ and $\not{D}_- = \delta_- + \overline{\delta}_-$ we see that

$$\overline{\partial_+^{\mathcal{N}}} = \overline{\partial}_+ + \overline{\mathcal{N}} \qquad d_- = \partial_- + \overline{\partial}_-.$$

And as a consequence of Corollary 3.4 and Propositions 7.3 and 7.4 we have

Proposition 7.8. *Let $(\mathcal{G}, \mathcal{I})$ be a positive SKT structure on a manifold with closed 3-form (M, H). Then the operators $\overline{\partial}_+^{\mathcal{N}}$ and d_- satisfy*

$$(\overline{\partial_+^{\mathcal{N}}})^2 = 0; \qquad \{\overline{\partial_+^{\mathcal{N}}}, d_-\} = 0; \qquad \{\partial_+^{\mathcal{N}}, \overline{\partial_+^{\mathcal{N}}}\} + d_-^2 = 0;$$

and

$$\overline{\partial_+^{\mathcal{N}}} : \Gamma(\wedge^k V_+^{0,1} \otimes \mathrm{Clif}(V_-)) \longrightarrow \Gamma(\wedge^{k+1} V_+^{0,1} \otimes \mathrm{Clif}(V_-));$$

$$d_- : \Gamma(\wedge^k V_+^{0,1}) \longrightarrow \Gamma(\wedge^k V_+^{0,1} \otimes V_-).$$

7.3 Representations

We finish our study of Lie algebroids and related objects associated to an SKT structure this section with some remarks about representations of the Lie algebroid $V_+^{1,0}$.

Firstly, it is worth pointing out that projection $\pi_T : V_+^{1,0} \longrightarrow TM$ has as its image $T^{1,0}M$, the $+i$-eigenspace of the complex structure introduced in Proposition 3.5 and hence $V_+^{1,0}$ is isomorphic to $T^{1,0}M$ as a Lie algebroid and a representation of one is the same as a representation of the other. Of course representations of $T^{1,0}M$ are simply holomorphic bundles endowed with a holomorphic connection (in fact, the connection determines the holomorphic structure), so one may just as well replace the words "representation of $V_+^{1,0}$" by "holomorphic vector bundle".

In Lemma 7.5 we saw that $V_-^{1,0}$ is a representation of $V_+^{1,0}$. In the proof we have used the fact that $\mathcal{J}_1$ is integrable, but that is not entirely necessary and in fact $V_-^{0,1}$ is also a representation of $V_+^{1,0}$.

Proposition 7.9. *Let $(M, \mathcal{G}, \mathcal{I})$ be a positive SKT manifold and $\mathcal{J}$ be a generalized complex extension of the SKT structure. The vector bundle $V_-^{0,1}$ is a representation of the Lie algebroid $V_+^{1,0}$ if endowed with the connection*

$$\nabla' : \Gamma(V_+^{1,0} \otimes V_-^{0,1}) \longrightarrow \Gamma(V_-^{0,1}); \qquad \nabla'_v w = \pi_{V_-^{0,1}}([\![v, w]\!]), \quad \forall v \in \Gamma(V_+^{1,0}), w \in \Gamma(V_-^{0,1}).$$

Proof. The proof that ∇' is a connection is the same as that given for ∇ in Lemma 7.5. So we only prove flatness, i.e.,

$$\nabla'_{v_1}\nabla'_{v_2} - \nabla'_{v_2}\nabla'_{v_1} - \nabla'_{[\![v_1,v_2]\!]} = 0 \qquad \text{for all } v_1, v_2 \in \Gamma(V_+^{1,0}).$$

Again, we first consider the term $\nabla'_{v_1}\nabla'_{v_2}w$. We decompose $[\![v_2,w]\!]$ into four components, according to the decomposition $\mathbb{T}_{\mathbb{C}}M = V_+^{1,0} \oplus V_+^{0,1} \oplus V_-^{1,0} \oplus V_-^{0,1}$. Since the failure of L_2 to be involutive with respect to the Courant bracket is given by the Nijenhuis tensor, $\overline{N} \in V_+^{0,1} \otimes \wedge^2 V_-^{1,0}$, we see that the component of $[\![v_2,w]\!]$ outside L_2 is given by $-\overline{N}(v_2,w) \in V_-^{1,0}$, so $[\![v_2,w]\!]$ has only three components and $\nabla'_{v_2}w$ is the component in $V_-^{0,1}$:

$$[\![v_2,w]\!] = u_+^{1,0} + \nabla'_{v_2}w - \overline{N}(v_2,w), \qquad u_+^{1,0} \in \Gamma(V_+^{1,0}).$$

Since $V_+^{1,0} \subset L_1$ and L_1 is integrable, we have that

$$[\![v_1, u_+^{1,0} - \overline{N}(v_2,w)]\!] \in L_1$$

hence the $V_-^{0,1}$ component of $[\![v_1, \nabla'_{v_2}w]\!]$ is the $V_-^{0,1}$ component of $[\![v_1,[\![v_2,w]\!]]\!]$, that is

$$\nabla'_{v_1}\nabla'_{v_2}w = \pi_{V_-^{0,1}}([\![v_1,[\![v_2,w]\!]]\!]).$$

Therefore we have

$$\nabla'_{v_1}\nabla'_{v_2}w - \nabla'_{v_2}\nabla'_{v_1}w - \nabla'_{[\![v_1,v_2]\!]}w = \pi_{V_-^{0,1}}([\![v_1,[\![v_2,w]\!]]\!] - [\![v_2,[\![v_1,w]\!]]\!] - [\![[\![v_1,v_2]\!],w]\!]) = 0,$$

where the last equality follows from the Jacobi identity. □

Finally, $V_- \otimes \mathbb{C} \cong T_{\mathbb{C}}M$ itself is a representation of $V_+^{1,0}$. Of course, $V_- \otimes \mathbb{C} = V_-^{1,0} \oplus V_-^{0,1}$ and since both summands are representations of $V_+^{1,0}$, so is the sum. Yet, the natural connection one should consider on $V_- \otimes \mathbb{C}$ is not the sum of the connections, but it should also include the Nijenhuis tensor of $\mathcal{J}_2$. This is also Bismut's "exotic holomorphic structure on $T_{\mathbb{C}}M$" [2].

Proposition 7.10. *Let $(M,\mathcal{G},\mathcal{I})$ be a positive SKT manifold. The vector bundle $V_{-\mathbb{C}} = V_- \otimes \mathbb{C}$ is a representation of the Lie algebroid $V_+^{1,0}$ if endowed with the connection*

$$\nabla^- : \Gamma(V_+^{1,0} \otimes V_{-\mathbb{C}}) \longrightarrow \Gamma(V_{-\mathbb{C}}); \qquad \widetilde{\nabla}_v w = \pi_{V_{-\mathbb{C}}}([\![v,w]\!]), \quad \textit{for all } v \in \Gamma(V_+^{1,0}), w \in \Gamma(V_{-\mathbb{C}}).$$

Further, if $\mathcal{J}$ is a generalized complex extension of the SKT structure, we can split $V_{-\mathbb{C}} = V_-^{1,0} \oplus V_-^{0,1}$ and in this splitting ∇^- is given by

$$\nabla^- = \begin{pmatrix} \nabla & -\overline{N} \\ 0 & \nabla' \end{pmatrix} \tag{7.8}$$

where ∇ is the connection introduced in Lemma 7.5 and ∇' is the connection introduced in Proposition 7.9

Proof. First we prove that in the presence of a generalized complex extension the definition of ∇^- agrees with the expression given in (7.8). Given $v \in \Gamma(V_+^{1,0})$, $w^{1,0} \in \Gamma(V_-^{1,0})$ and $w^{0,1} \in \Gamma(V_-^{0,1})$ we have that $[\![v, w^{1,0}]\!]$ lies in L_1 by integrability of $\mathcal{J}_1$ hence its V_- component is its $V_-^{1,0}$ component which is, by definition, $\nabla_v w^{1,0}$.

Similarly, $[\![v, w^{0,1}]\!]$ has three components, as established in the previous Proposition:

$$[\![v, w^{0,1}]\!] = u_+ + \nabla'_v w^{0,1} - \overline{N}(v, w^{0,1}),$$

hence, projecting onto V_- we get

$$\nabla^-_v w^{0,1} = \nabla'_v w^{0,1} - \overline{N}(v, w^{0,1})$$

and hence ∇^- is given by (7.8)

Further, similarly to the previous case, we have that for $v_1, v_2 \in \Gamma(V_+^{1,0})$

$$[\![v_2, w^{1,0}+w^{0,1}]\!] = u_+^{1,0}+\nabla_v w^{1,0}+\nabla'_v w^{0,1}-N(v, w^{0,1}) = u_+^{1,0}+\nabla^-_v(w^{1,0}+w^{0,1}), \quad u_+^{1,0} \in \Gamma(V_+^{1,0}).$$

And again, since $V_+^{1,0}$ is involutive, we see that

$$\nabla^-_{v_1}\nabla^-_{v_2} w = \pi_{V_-}([\![v_1, \pi_{V_-}([\![v_2, w]\!])]\!]) = \pi_{V_-}([\![v_1, [\![v_2, w]\!]]\!]),$$

and flatness of the connection follows once again from the Jacobi identity for the Courant bracket. □

With these representations at hand, one can also see the operator $\overline{\partial}_+$ as studied in Proposition 7.4 as an extension of the Lie algebroid differential $\overline{\partial}_+ : \Gamma(\wedge^k V_+^{0,1}) \longrightarrow \Gamma(\wedge^{k+1} V_+^{0,1})$ to take values on the tensor product of the exterior algebra of the flat modules $V_-^{1,0}$ and $V_-^{0,1}$. Similarly, the operator $\overline{\partial_+^N}$ is the extension of the Lie algebroid differential $\overline{\partial}_+$ to take values in the representation from Proposition 7.10.

8 Deformations

One can consider at least three different deformation problems. In the most strict of them one fixes the generalized metric and asks for deformations of the complex structure on V_+. For the second deformation problem, one fixes the Courant algebroid where the SKT structure is defined and looks for deformations of metric and complex structure which are still SKT. In the third, and more general problem, one lets the Courant algebroid change as well, that is, if ω_t is the 1-parameter family of Hermitian forms, one only requires that $dd^c_t\omega_t = 0$, but the cohomology class $[d^c_t\omega_t]$ is allowed to vary with t.

Here we will consider the first two problems. In both cases, we want to find involutive deformations, $V_+^{1,0\varepsilon}$, of $V_+^{1,0}$, in the first problem we require $V_+^{1,0\varepsilon}$ to lie in V_+, while in the second problem that condition is removed. Either way, as long as the deformation is small enough, the remaining SKT conditions, namely,

$$V_+^{1,0\varepsilon} \cap \overline{V_+^{1,0\varepsilon}} = \{0\}$$

and

$$\langle v, v\rangle > 0 \qquad \text{for all } v \in V_+^{1,0\varepsilon} \oplus \overline{V_+^{1,0\varepsilon}}, v = \overline{v}$$

will still hold, since these are open conditions and hence if $V_+^{1,0\varepsilon}$ is involutive, it defines an SKT structure.

8.1 First deformation problem

The first problem is very similar to the problem of deformations of generalized complex structures. Indeed, since V_+ is to remain fixed, any small deformation of $V_+^{1,0}$ is still transversal to $V_+^{0,1}$ and hence can be described by the graph of a bundle map $\varepsilon : V_+^{1,0} \longrightarrow V_+^{0,1}$:

$$V_+^{1,0\varepsilon} := \{v^{1,0} + \varepsilon(v^{1,0}) : v^{1,0} \in V_+^{1,0}\}.$$

Since $V_+^{0,1} \cong (V_+^{1,0})^*$, ε is a section of $\otimes^2 V_-^{0,1}$. Since $V_+^{1,0\varepsilon}$ is isotropic, we have that $\varepsilon \in \Gamma(\wedge^2 V_+^{0,1})1$.

Theorem 8.1. *Let (M, H) be a manifold with 3-form and $(\mathcal{G}, \mathcal{I})$ be a positive SKT structure on (M, H). An element $\varepsilon \in \Gamma(\wedge^2 V_+^{0,1})$ of small norm gives rise to an integrable SKT structure $\mathcal{I}_\varepsilon$ compatible with the metric $\mathcal{G}$ if and only if*

$$\begin{aligned} d_-\varepsilon &= 0; \\ \overline{\partial^{\mathcal{N}}_+}\varepsilon + \frac{1}{2}[\![\varepsilon, \varepsilon]\!]_{SN} &= 0, \end{aligned} \tag{8.1}$$

where $[\![\cdot, \cdot]\!]_{SN}$ is the Schouten–Nijenhuis bracket extending the Courant bracket to $\Gamma(\wedge^\bullet V_+^{0,1})$.

If $\mathcal{J}$ is a generalized complex extension of the SKT structure, the equations above become

$$\partial_-\varepsilon = \overline{\partial}_-\varepsilon = 0$$

$$\overline{\partial}_+\varepsilon + \frac{1}{2}[\![\varepsilon, \varepsilon]\!]_{SN} = 0.$$

Proof. For the deformation given by $\varepsilon \in \Gamma(\wedge^2 V_+^{0,1}) \subset \mathrm{Clif}(\mathbb{T}_{\mathbb{C}}M)$ as above, the space of forms which annihilate $V_+^{1,0\varepsilon}$, $\mathcal{W}_\varepsilon^n$, is given by $\mathcal{W}_\varepsilon^n = e^{-\varepsilon} \cdot \mathcal{W}^n$ and integrability is equivalent to

$$d^H : \mathcal{W}_\varepsilon^n \longrightarrow \mathcal{W}_\varepsilon^n \oplus \mathcal{W}_\varepsilon^{n-2}.$$

But, since V_+ and V_- do not change under the deformation, elements in $\mathcal{W}_\varepsilon^n \oplus \mathcal{W}_\varepsilon^{n-2}$ are those which can be obtained from $\mathcal{W}_\varepsilon^n$ by the Clifford action of elements in $V_+ \otimes \mathrm{Clif}(V_-)$ and $\mathcal{W}_\varepsilon^n = e^{-\varepsilon} \cdot \mathcal{W}^n$, so the condition above holds if and only if for all $\varphi \in \mathcal{W}^n$ there is $\alpha \in V_+ \otimes \mathrm{Clif}(V_-)$ such that

$$d^H(e^{-\varepsilon} \cdot \varphi) = \alpha \cdot e^{-\varepsilon} \cdot \varphi = e^{-\varepsilon} \cdot ((e_*^\varepsilon \alpha) \cdot \varphi).$$

And since $\varepsilon \in \wedge^2 V_+^{0,1}$ $e_*^\varepsilon \alpha \in V_+ \otimes \mathrm{Clif}(V_-)$, as e^ε only acts in the V_+ part as a complex orthogonal transformation. Therefore, integrability of the deformed structure is equivalent to

$$e^\varepsilon d^H e^{-\varepsilon}(\varphi) = \widetilde{\alpha} \cdot \varphi \qquad \text{for all } \varphi \in \mathcal{W}^n, \text{ for some } \widetilde{\alpha} \in \Gamma(V_+ \otimes \mathrm{Clif}(V_-)).$$

Now we can expand the operator on the left as a series:

$$e^\varepsilon d^H e^{-\varepsilon}(\varphi) = d^H + [\varepsilon, d^H] + \frac{1}{2!}[\varepsilon, [\varepsilon, d^H]] + \frac{1}{3!}[\varepsilon, [\varepsilon, [\varepsilon, d^H]]] + \cdots$$

The third term of the series is computed in terms of the derived bracket of ε, with itself, that is, the extension of the Schouten–Nijenhuis bracket to sections of the Lie algebroid $V_+^{0,1}$:

$$[\varepsilon, [\varepsilon, d^H]] = \{\varepsilon, \{\varepsilon, d^H\}\} = -\{\{\varepsilon, d^H\}, \varepsilon\} = -[\![\varepsilon, \varepsilon]\!]_{SN} \in \Gamma(\wedge^3 V_+^{0,1}).$$

Since $V_+^{0,1}$ is isotropic, all terms of the series after the third vanish and we have

$$e^{\varepsilon} d^H e^{-\varepsilon}(\varphi) = (d^H + [\varepsilon, d^H] - \frac{1}{2}[\![\varepsilon, \varepsilon]\!]_{SN})\varphi.$$

Since the original SKT structure is integrable, we have that $d^H : \mathcal{W}^n \longrightarrow \mathcal{W}^n \oplus \mathcal{W}^{n-2}$. Splitting $d^H = \not{D}_-^H + \delta_+^N + \overline{\delta_+^N}$ we have that $[\varepsilon, \delta_+^N]$ maps $\mathcal{W}^n$ into $\mathcal{W}^{n-2}$. So integrability of the deformed structure is equivalent to the requirement that the remaining terms also map $\mathcal{W}^n$ into $\mathcal{W}^n \oplus \mathcal{W}^{n-2}$.

Now notice that $[\varepsilon, \not{D}_-^H] : \mathcal{W}^n \longrightarrow \mathcal{W}^{n-4}$ while $[\varepsilon, \delta_+^N] - \frac{1}{2}[\![\varepsilon, \varepsilon]\!]_{SN} : \mathcal{W}^n \longrightarrow \mathcal{W}^{n-6}$, so for the deformed structure to be integrable, each of these maps must vanish. From Proposition 7.8, these conditions are equivalent to the conditions presented in the statement of the theorem. □

8.2 Second deformation problem and stability

As we mentioned earlier, in the second deformation problem, one fixes the Courant algebroid and looks for deformations of the SKT structure. Stated this way, the problem has an infinite dimensional space of solutions even after taking the quotient by the action of the diffeomorphism group. Indeed, from the classical viewpoint, an SKT structure is a Hermitian structure (g, I) for which $dd^c\omega = 0$. For any $f \in C^\infty(M)$ we can consider a new Hermitian form: $\omega + \varepsilon dd^c f$. In a compact manifold if we take ε small enough this gives rise to new non diffeomorphic SKT structures on M still with 3-form $d^c\omega$. In fact, more can be said about the set of deformations of an SKT structure which preserve the complex structure I. The author has learnt from Swann and Zabzine a small variation of the following proposition, which can also be found, rephrased, in [11]:

Proposition 8.2. *The set of small deformations of an SKT structure (g, I) which preserve the complex structure and the cohomology class $[d^c\omega]$ is parametrized by the kernel of the map*

$$d^c : \Omega^{1,1}(M) \longrightarrow \Omega^3(M)/\Omega^3_{exact}(M).$$

In particular there is a disc around the zero form in the set of closed 2-forms on M with nontrivial $(1,1)$ component, $D \subset \Omega^2_{cl}(M)/(\Omega^{2,0}_{cl}(M) + \Omega^{0,2}_{cl}(M))$ which maps injectively into the kernel of the map above.

Proof. The first claim is clear: a variation of the metric is equivalent to a deformation of the Hermitian form $\omega = g \circ I$ and since I is orthogonal with respect to the deformed structure, ω must change by a form $B \in \Omega^{1,1}(M)$. Then the SKT condition gives

$$d^c(\omega + B) = H + d^c B,$$

and since this is to have the same cohomology class as H, $d^c B$ must be exact.

Now, given any closed 2-form B, let $B^{p,q}$ be its component in $\Omega^{p,q}(M)$. Then let $\widetilde{\omega} = \omega + B^{1,1}$. As long as $B^{1,1}$ is small, nondegeneracy and positivity of $g = -\widetilde{\omega} \circ I$ still hold as those are open conditions. And finally,

$$\begin{aligned} d^c(\omega + B^{1,1}) &= H - i\partial B^{1,1} + i\overline{\partial} B^{1,1} = H + i\overline{\partial} B^{2,0} - i\partial B^{0,2} \\ &= H + d(iB^{2,0} - iB^{0,2}) = H + 2d\mathrm{Im}(B^{2,0}). \end{aligned}$$

□

The fact that one can so freely deform the metric and remain in the SKT category is a very useful phenomenon which, for example, allows one to blow-up complex submanifolds of SKT manifolds [13]. Yet, from the point of view of deformations, these seem a little less interesting than deformations which change the complex structure.

Therefore it is natural to phrase the second deformation problem as a stability problem. In our setup, the direct translation of the classical question is as follows: Given an SKT structure $(\mathcal{G}, \mathcal{I})$, Proposition 3.5 gives us an extension to a generalized Hermitian structure $(\mathcal{G}, \mathcal{J}_1)$ for which $\mathcal{J}_1$ is of complex type. Then one is interested in which deformations of $\mathcal{J}_1$ as a generalized complex structure of complex type give rise to deformations of the SKT structure.

What we do next is phrased in slightly more general terms in order to accommodate other cases of interest. Our proof is inspired on the approach used by Goto to prove the generalized Kähler stability theorem [16].

Theorem 8.3. (SKT Stability Theorem) *Let $(\mathcal{G}, \mathcal{J}_1)$ be a generalized Hermitian extension of a positive SKT structure. Given $\mathcal{J}_{1\varepsilon}$, an analytic family of deformations of $\mathcal{J}_1$ parametrized by a small disc $D \subset \mathbb{C}$, $\varepsilon : D \longrightarrow \Gamma(\wedge^2 \overline{L_1})$, we let*

$$\begin{aligned} \varepsilon^+ &: D \longrightarrow \Gamma(\wedge^2 V_+^{0,1}) \\ \varepsilon^\pm &: D \longrightarrow \Gamma(V_+^{0,1} \otimes V_-^{0,1}) \\ \varepsilon^- &: D \longrightarrow \Gamma(\wedge^2 V_-^{0,1}) \end{aligned}$$

be the components of ε and for each of them let $\varepsilon^\bullet = \sum_{k=1}^\infty \frac{1}{k!}\varepsilon_k^\bullet t^k$ be the corresponding series expansion. Then the obstructions to finding a generalized metric $\mathcal{G}_\varepsilon$ which makes $(\mathcal{G}_\varepsilon, \mathcal{J}_{1\varepsilon})$ into a generalized Hermitian extension of an SKT structure lie in the space $H^{2,1}_{\overline{\partial}_+}(M; \overline{L_2})$ and the first obstruction is the class $[\partial_- \varepsilon_1^+ + \overline{\mathcal{N}} \varepsilon_1^\pm]$.

According to the deformation theorem for Lie bialgebroids [21, 27], a deformation of $\mathcal{J}_1$ is given by sections $\varepsilon(t)$ as above, which we will denote simply by ε, satisfying the Maurer–Cartan equation and the corresponding deformed structure is given by

$$\mathcal{J}_{1\varepsilon} = I_\varepsilon \mathcal{J}_1 I_\varepsilon^{-1},$$

where I_ε is the real transformation which in the splitting $\mathbb{T}_{\mathbb{C}} M = L_1 \oplus \overline{L_1}$ is given by the matrix

$$I_\varepsilon = \begin{pmatrix} \mathrm{Id} & \overline{\varepsilon} \\ \varepsilon & \mathrm{Id} \end{pmatrix}.$$

Since this matrix is not orthogonal with respect to the natural pairing, in general, neither is $I_\varepsilon \mathcal{J}_2 I_\varepsilon^{-1}$ and hence a complex structure on $\mathbb{T}M$ defined this way is not even an almost generalized complex structure. The first step is to translate this problem into deformations using orthogonal transformations so that we can find an generalized almost complex structure $\mathcal{J}_{2\varepsilon}$ making $\mathcal{J}_{1\varepsilon}$ into a generalized Hermitian structure.

Lemma 8.4. There is a map $a : D \subset \wedge^2\overline{L_1} \longrightarrow \wedge^2(L_1 \oplus \wedge^2\overline{L_1})_{\mathbb{R}}$, where D is a small disc around the origin and $(\cdot)_{\mathbb{R}}$ denotes the real elements in the vector space, such that for each $\varepsilon \in \Gamma(\wedge^2\overline{L_1})$, $a(\varepsilon)$ is the unique element in the codomain that satisfies

$$e_*^a \mathcal{J}_1 e_*^{-a} = I_\varepsilon \mathcal{J}_1 I_\varepsilon^{-1}.$$

Further $a_*|_0(\varepsilon) = -(\varepsilon + \overline{\varepsilon})$.

Proof. Indeed, the space of generalized complex structures of the same parity as $\mathcal{J}_1$ on T_pM is the homogeneous space $SO(\mathbb{T}_pM)/\mathrm{Stab}(\mathcal{J}_1) \cong SO(n,n)/U(n,n)$. Hence, composing the exponential map with the projection

$$\mathfrak{so}(\mathbb{T}_pM) \longrightarrow SO(\mathbb{T}_pM) \longrightarrow SO(\mathbb{T}_pM)/\mathrm{Stab}(\mathcal{J}_1)$$

gives a local submersion. Since the elements in $\mathfrak{so}(\mathbb{T}_pM)$ preserving $\mathcal{J}_1$ are those in $(L_1 \otimes \overline{L_1})_{\mathbb{R}}$, and $(\wedge^2 L_1 \oplus \wedge^2\overline{L_1})_{\mathbb{R}}$ is a complementary subspace, we have that

$$(\wedge^2 L_1 \oplus \wedge^2\overline{L_1})_{\mathbb{R}} \longrightarrow SO(\mathbb{T}_pM)/\mathrm{Stab}(\mathcal{J}_1).$$

is a local diffeomorphism. That is, for each small deformation of a generalized complex structure $\mathcal{J}_1$ on V there is a unique element $a \in (\wedge^2 L_1 \oplus \wedge^2\overline{L_1})_{\mathbb{R}}$ which realizes it.

For the last claim, we observe that if $\varepsilon : (-s,s) \longrightarrow \wedge^2\overline{L_1}$ is a path which passes through 0 at time zero and $v \in L_1$, then $\dot{I}_\varepsilon|_0(v+\overline{v}) = \dot{\varepsilon}|_0(v) + \dot{\overline{\varepsilon}}|_0(\overline{v})$. On the other hand, for $a : (-s,s) \longrightarrow \in (\wedge^2 L_1 \oplus \wedge^2\overline{L_1})_{\mathbb{R}}$, say $a = \alpha + \overline{\alpha}$ with $\alpha \in \wedge^2 L_1$, we have

$$\tfrac{d}{dt} e_*^a|_0(v+\overline{v}) = \tfrac{d}{dt}(e^a(v+\overline{v})e^{-a})|_0 = \{a|_0, v+\overline{v}\} = -(\alpha|_0(v) + \overline{\alpha}|_0(\overline{v})),$$

hence if the family a gives rise to the same deformation as ε, we must have $a = -(\varepsilon + \overline{\varepsilon})$. ☐

Proof of Theorem 8.3. With this setup, e_*^a is an orthogonal transformation of $\mathbb{T}M$ and hence $e_*^a \mathcal{J}_2 e_*^{-a}$ is an generalized almost complex structure which commutes with $\mathcal{J}_{1\varepsilon}$. If $b \in \mathbb{R} \oplus (L_1 \otimes \overline{L_1})_{\mathbb{R}} \subset \mathrm{Clif}^2(\mathbb{T}M)$ the orthogonal transformation e_*^b preserves $\mathcal{J}_1$ hence $\mathcal{J}_{1\varepsilon}$ and $\mathcal{J}_{2\varepsilon} = e_*^a e_*^b \mathcal{J}_2 e_*^{-b} e_*^{-a}$ give rise to a generalized Hermitian structure for any choice of b. Our task is to identify suitable conditions on a given a under which we can choose b so that the pair $\mathcal{J}_{1\varepsilon}$ and $\mathcal{J}_{2\varepsilon}$ gives an SKT structure.

Since $\mathcal{J}_{1\varepsilon}$ is integrable, we know that

$$d^H : \mathcal{U}_\varepsilon^{0,n} \longrightarrow \mathcal{U}_\varepsilon^{1,n-1} \oplus \mathcal{U}_\varepsilon^{1,n-3} \oplus \mathcal{U}_\varepsilon^{-1,n-1} \oplus \mathcal{U}_\varepsilon^{-1,n-3}$$

and, according to Theorem 3.6, the SKT condition is that the $\mathcal{U}_\varepsilon^{-1,n-3}$ component of the map above must vanish.

Since $\mathcal{J}_{1\varepsilon}$ and $\mathcal{J}_{2\varepsilon}$ are obtained from $\mathcal{J}_1$ and $\mathcal{J}_2$ by the action of $e^a e^b \in Spin(\mathbb{T}M)$, the corresponding decompositions of spinors are related by

$$\mathcal{U}^{p,q}_{\varepsilon} = e^a e^b \mathcal{U}^{p,q},$$

so the different components of d^H in the splitting induced by $(\mathcal{J}_{1\varepsilon}, \mathcal{J}_{2\varepsilon})$ (c.f. Figure 3)

$$d^H = \delta_{+\varepsilon} + \overline{\delta}_{+\varepsilon} + \delta_{-\varepsilon} + \overline{\delta}_{-\varepsilon} + N_{\varepsilon} + \overline{N_{\varepsilon}} + N_{2\varepsilon} + \overline{N_{2\varepsilon}}$$

are in correspondence with the components of $e^{-b}e^{-a}d^H e^a e^b$ in the splitting determined by $(\mathcal{J}_1, \mathcal{J}_2)$ (see Figure 12).

$$\begin{tikzcd}
 & & \mathcal{U}^{0,n} \arrow[dl, swap, "e^{-a}e^{-b}\overline{\delta}_{+\varepsilon}e^a e^b"] \arrow[dr, "e^{-a}e^{-b}\delta_{-\varepsilon}e^a e^b"] \arrow[dddl, swap, "e^{-a}e^{-b}\overline{N_{2\varepsilon}}e^a e^b"] \arrow[dddr, "e^{-a}e^{-b}\overline{N_{\varepsilon}}e^a e^b"] & & \\
 & \mathcal{U}^{-1,n-1} & & \mathcal{U}^{1,n-1} & \\
 & & \mathcal{U}^{0,n-2} & & \\
\iddots & \mathcal{U}^{-1,n-3} & & \mathcal{U}^{1,n-3} & \ddots
\end{tikzcd}$$

Figure 12: Nontrivial components of $e^{-b}e^{-a}d^H e^a e^b$ and their relation to the components of d^H with respect to the decomposition of forms determined by $(\mathcal{J}_{1\varepsilon}, \mathcal{J}_{2\varepsilon})$.

Since $a \in (\wedge^2 L_1 \oplus \wedge^2 \overline{L_1})_{\mathbb{R}}$, there are $\alpha^+ \in \Gamma(\wedge^2 V_+^{0,1})$, $\alpha^{\pm} \in \Gamma(V_+^{0,1} \otimes V_-^{0,1})$ and $\alpha^- \in \Gamma(\wedge^2 V_-^{0,1})$ such that

$$a = \alpha^+ + \alpha^{\pm} + \alpha^- + \overline{\alpha^+} + \overline{\alpha^{\pm}} + \overline{\alpha^-}. \tag{8.2}$$

Similarly, any $b \in \Gamma(\mathbb{R} \oplus L_1 \otimes \overline{L_1})_{\mathbb{R}}$ can be decomposed in four components according to the splitting

$$\mathbb{R} \oplus (L_1 \otimes \overline{L_1})_{\mathbb{R}} = \mathbb{R} \oplus (V_+^{1,0} \otimes V_+^{0,1})_{\mathbb{R}} \oplus (V_-^{1,0} \otimes V_-^{0,1})_{\mathbb{R}} \oplus ((V_+^{0,1} \otimes V_-^{1,0}) \oplus (V_+^{1,0} \otimes V_-^{0,1})_{\mathbb{R}})$$

however, as we will see later, only the last of these components is relevant for the deformation problem. So we let b is of the form

$$b = \beta + \beta^{\pm} + \overline{\beta^{\pm}} \tag{8.3}$$

with $\beta^{\pm} \in \Gamma(V_+^{0,1} \otimes V_-^{1,0})$ and $\beta \in \Gamma(\mathbb{R} \oplus (V_+^{1,0} \otimes V_+^{0,1})_{\mathbb{R}} \oplus (V_-^{1,0} \otimes V_-^{0,1})_{\mathbb{R}})$.

Since the deformation family $\varepsilon(t)$ is analytic on t, so is $a(t)$ and we can try to solve the condition that the $\mathcal{U}^{-1,n-3}$ component of $e^{-b}e^{-a}d^H e^a e^b \psi$ vanishes by a power series argument. In the sequence given an analytic function $f(t)$, where f can be α^+, $\alpha^{\pm}$ and so on, we denote denote the coefficients of its power series expansion by f_k, so that $f = \sum \frac{1}{k!} f_k t^k$.

The first nontrivial condition on a imposed by the integrability of $\mathcal{J}_1$ is that $[d^H, a_1]$ must have no components in $\mathcal{U}^{n-3,\bullet}$. In terms of the decomposition of a above, this implies

$$[\overline{\delta}_+, \alpha_1^+] = 0 \qquad [\overline{\delta}_+, \alpha_1^\pm] + [\overline{\delta}_-, \alpha_1^+] = 0$$

or equivalently

$$\overline{\partial}_+\alpha_1^+ = 0 \qquad \overline{\partial}_+\alpha_1^\pm + \overline{\partial}_-\alpha_1^+ = 0. \tag{8.4}$$

The first term in the series expansion of $e^{-b}e^{-a}d^He^ae^b\psi$ is

$$[d^H, b_1 + a_1]\psi.$$

Splitting a and b into their components, we see that the requirement that the $\mathcal{U}^{-1,n-3}$ component vanishes is equivalent to

$$([\overline{\delta}_+, \beta_1^\pm] + [\delta_-, \alpha_1^+] + [\overline{N}, \alpha_1^\pm])\psi = 0$$

Isolating $\beta^\pm$ and using that ψ is nonzero, we can rewrite the condition above as

$$\overline{\partial}_+\beta_1^\pm = -\partial_-\alpha_1^+ - \overline{\mathcal{N}}\alpha_1^\pm.$$

And one obvious necessary condition for this to have a solution is that the right hand side must be $\overline{\partial}_+$-closed. That is indeed the case, since

$$\overline{\partial}_+(\partial_-\alpha_1^+ + \overline{\mathcal{N}}\alpha_1^\pm) = -\partial_-\overline{\partial}_+\alpha_1^+ - \overline{\partial}_-\overline{\mathcal{N}}\alpha_1^+ - \overline{\mathcal{N}}\overline{\partial}_-\alpha_1^+ - \overline{\mathcal{N}}\overline{\partial}_+\alpha_1^\pm = 0$$

where for the first equality we used the commutation rule for $\overline{\partial}_+$ and ∂_- from Proposition 7.4 and in the second equality we used equations (8.4) and the fact that $\mathcal{N}\alpha_1^+ = 0$ as $\alpha_1^+ \in \overline{L_2}$. If this $\overline{\partial}_+$-closed form is in fact $\overline{\partial}_+$-exact, we can use $G_{\overline{\partial}_+}$, the Green operator for $\overline{\partial}_+$, to find a suitable $\beta_1^\pm$:

$$\beta_1^\pm = -\overline{\partial}_+^* G_{\overline{\partial}_+}(\partial_-\alpha_1^+ + \overline{\mathcal{N}}\alpha_1^\pm)$$

Finally, we recall that according to Lemma 8.4, $a_1 = -(\varepsilon_1 + \overline{\varepsilon_1})$, so the components of a_1 are determined by the different components of ε_1 hence the equation $\beta_1^\pm$ must solve is

$$\overline{\partial}_+\beta_1^\pm = \partial_-\varepsilon_1^+ + \overline{\mathcal{N}}\varepsilon_1^\pm,$$

so the first obstruction class is $[\partial_-\varepsilon_1^+ + \overline{\mathcal{N}}\varepsilon_1^\pm] \in H^{2,1}_{\overline{\partial}_+}(M; \overline{L_2})$.

Now we move to the general case, which is proved by induction. We assume that we have chosen a_i and b_i for $i < k$ such that the component of $e^{-b}e^{-a}d^He^ae^b$ mapping $\mathcal{U}^{0,n}$ into $\mathcal{U}^{-1,n-3}$ vanishes to order $k-1$. Then the vanishing of the order k component of this map is the condition

$$[\beta_k^\pm, \overline{\delta}_+] + [\alpha_k^+, \delta_-] + [\alpha_k^\pm, \overline{N}] + F_k(a_1, \cdots, a_{k-1}, b_1, \cdots, b_{k-1}) = 0,$$

where F_k is some function. This condition is equivalent to

$$\overline{\partial_+}\beta_k^\pm = -(\partial_-\alpha_k^+ + \overline{\mathcal{N}}\alpha_k^\pm + F_k(a_1, \cdots, a_{k-1}, b_1, \cdots, b_{k-1})).$$

And if there is $\beta_k^\pm$ which solves this equation, we can proceed to the next step. If for all possible choices of b_i for $i<k$ which guarantee the vanishing of the lower order terms this equation has no solution, then the deformation is obstructed.

Next we show that $\partial_-\alpha_k^+ + \overline{\mathcal{N}}\alpha_k^\pm + F_k(a_1,\cdots,a_{k-1},b_1,\cdots,b_{k-1})$ is $\overline{\partial}_+$-closed so we can conclude that the obstruction space is $H^{2,1}(M;\overline{L_2})$. Since $\overline{\partial}_+\beta_k^\pm$ is clearly $\overline{\partial}_+$-closed, we must prove that

$$\overline{\partial}_+(\overline{\partial}_+\beta_k^\pm + \partial_-\alpha_k^+ + \overline{\mathcal{N}}\alpha_k^\pm + F_k(a_1,\cdots,a_{k-1},b_1,\cdots,b_{k-1})) = 0$$

but the term in parenthesis is precisely the order k component of $e^{-b}e^{-a}\overline{N_{2\varepsilon}}e^a e^b$ (see Figure 12) and by assumption, the lower order terms of $e^{-b}e^{-a}\overline{N_{2\varepsilon}}e^a e^b$ vanish, therefore we must prove that

$$\overline{\partial}_+ e^{-b}e^{-a}\overline{N_{2\varepsilon}}e^a e^b = \{\overline{\delta}_+, e^{-b}e^{-a}\overline{N_{2\varepsilon}}e^a e^b\} \tag{8.5}$$

vanishes to order k. But since $\overline{\delta}_+ = e^{-b}e^{-a}\overline{\delta}_{+\varepsilon}e^b e^a + O(t)$, we see that (8.5) vanishes to order k if and only if

$$\{e^{-b}e^{-a}\overline{\delta}_{+\varepsilon}e^a e^b, e^{-b}e^{-a}\overline{N_{2\varepsilon}}e^a e^b\}$$

vanishes to order k. But for this operator, we have

$$\{e^{-b}e^{-a}\overline{\delta}_{+\varepsilon}e^a e^b, e^{-b}e^{-a}\overline{N_{2\varepsilon}}e^a e^b\} = e^{-b}e^{-a}\{\delta_{+\varepsilon}, \overline{N_{2\varepsilon}}\}e^a e^b = 0,$$

where in the last equality we used that, in a generalized Hermitian manifold, the condition $(d^H)^2 = 0$ gives, among other things, that $\{\overline{\delta}_+, \overline{N_2}\} = 0$.

Finally, if all obstructions vanish, standard elliptic estimates show that the sequence constructed above converges. See for example [16] for the proof of convergence for the analogous problem in the generalized Kähler case. □

It is interesting to notice that one can deform the generalized complex extension $\mathcal{J}_1$ without deforming the SKT structure.

Proposition 8.5. *Let* $(\mathcal{G},\mathcal{J}_1)$ *be a generalized complex extension of a positive SKT structure* $(\mathcal{G},\mathcal{I})$ *on a manifold* M. *Let* $\varepsilon \in \Gamma(\wedge^2 V_-^{0,1}) \subset \Gamma(\wedge^2\overline{L_1})$ *be a Maurer–Cartan element, i.e., the corresponding deformation of* $\mathcal{J}_1$, $\mathcal{J}_{1\varepsilon}$, *is integrable. Then, if* ε *is small enough,* $(\mathcal{G},\mathcal{J}_{1\varepsilon})$ *is a generalized complex extension of the SKT structure* $(\mathcal{G},\mathcal{I})$.

Proof. According to Lemma 8.4, if ε is small enough, there is an element $a \in \Gamma(\wedge^2 L_1 \oplus \wedge^2\overline{L_1})_{\mathbb{R}}$ such that

$$e_*^{-a}\mathcal{J}_1 e_*^a = e_*^{-\varepsilon}\mathcal{J}_1 e^\varepsilon$$

but since $\varepsilon \in \wedge^2 V_-^{0,1}$ one can easily see that $a \in \Gamma(\wedge^2 V_-^{1,0} \oplus \wedge^2 V_-^{0,1})$. Then $e^a|_{V_+} = \mathrm{Id}$ and $e^a|_{V_-}$ is an orthogonal transformation of V_-, therefore $e^{-a}\mathcal{G}e^a = \mathcal{G}$ and $e^{-a}\mathcal{J}_1 e^a|_{V_+} = \mathcal{J}_1|_{V_+} = \mathcal{I}$. That is $e^{-a}\mathcal{J}_1 e^a$ is a generalized complex extension of $(\mathcal{G},\mathcal{I})$. □

Of course, the same arguments with the obvious changes give deformation results for negative SKT structures (c.f. Figure 8):

Theorem 8.6. (Negative SKT Stability Theorem) *Let $(\mathcal{G}, \mathcal{J}_1)$ be a generalized Hermitian extension of a negative SKT structure. Given $\mathcal{J}_{1\varepsilon}$, an analytic family of deformations of $\mathcal{J}_1$ parametrized by a small disc $D \subset \mathbb{C}$, $\varepsilon : D \longrightarrow \Gamma(\wedge^2\overline{L_1})$, we let*

$$\begin{aligned}\varepsilon^+ &: D \longrightarrow \Gamma(\wedge^2 V_+^{0,1})\\ \varepsilon^\pm &: D \longrightarrow \Gamma(V_+^{0,1} \otimes V_-^{0,1})\\ \varepsilon^- &: D \longrightarrow \Gamma(\wedge^2 V_-^{0,1})\end{aligned}$$

be the components of ε and for each of them let $\varepsilon^\bullet = \sum_{k=1}^{\infty} \varepsilon_k^\bullet t^k$ be the corresponding series expansion. Then the obstructions to finding a generalized metric $\mathcal{G}_\varepsilon$ which makes $(\mathcal{G}_\varepsilon, \mathcal{J}_{1\varepsilon})$ into a generalized Hermitian extension of a negative SKT structure lie in the space $H^{2,1}_{\partial_-}(M; \overline{L_2})$ and the first obstruction is the class $[\overline{\partial}_+\overline{\varepsilon_1^-} + \overline{\mathcal{N}}_2\overline{\varepsilon_1^{\pm}}]$.

Proposition 8.7. *Let $(\mathcal{G}, \mathcal{J}_1)$ be a generalized complex extension of a negative SKT structure $(\mathcal{G}, \mathcal{I})$ on a manifold M. Let $\varepsilon \in \Gamma(\wedge^2 V_+^{0,1})$ be a Maurer–Cartan element and $\mathcal{J}_{!\varepsilon}$ be the corresponding deformation of $\mathcal{J}_1$. Then, if ε is small enough, $(\mathcal{G}, \mathcal{J}_{1\varepsilon})$ is a generalized complex extension of the SKT structure $(\mathcal{G}, \mathcal{I})$.*

8.3 Deformations of generalized Kähler structures

Similarly to the SKT deformation problem studied in Section 8.2, a common way to study the question of deformations of a (generalized) Kähler structure $(\mathcal{J}_1, \mathcal{J}_2)$ is in the context of stability: For which deformations of $\mathcal{J}_1$ is there a corresponding deformation of $\mathcal{J}_2$ such that the pair of deformed structures is still a (generalized) Kähler structure?

In its classical setting, this question was successfully answered by Kodaira [24], where $\mathcal{J}_1$ is taken to be the complex structure and in the generalized setting, for analytic deformations, by Goto [16] under the additional hypothesis that the canonical bundle of $\mathcal{J}_2$ has a globally defined nowhere vanishing closed section. In both cases, there are no obstructions to the problem. However, even in the Kähler setting, if we let $\mathcal{J}_1$ be the symplectic structure and $\mathcal{J}_2$ be the complex, the problem does have obstructions [8].

As we have seen in Section 2, a generalized Kähler structure is at the same time a positive and a negative SKT structure. Hence, when studying the stability problem of a generalized Kähler structure there are three possible outcomes one can expect:

1. The deformation is unobstructed and one can complete the deformed $\mathcal{J}_1$ to a generalized Kähler structure;
2. The deformation is half obstructed and one can complete the deformed $\mathcal{J}_1$ to a positive or negative SKT structure;
3. The deformation is fully obstructed and $\mathcal{J}_1$ can not be completed to either a positive or a negative SKT structure.

Theorem 8.8 (Generalized Kähler Stability Theorem)**.** *Let $(M, \mathcal{J}_1, \mathcal{J}_2)$ be a compact generalized Kähler manifold. Given $\mathcal{J}_{1\varepsilon}$, an analytic family of deformations of $\mathcal{J}_1$ parametrized* *by a small disc* $D \subset \mathbb{C}$, $\varepsilon : D \longrightarrow \Gamma(\wedge^2 \overline{L_1})$, *we let*

$$\begin{aligned}
\varepsilon^+ &: D \longrightarrow \Gamma(\wedge^2 V_+^{0,1}) \\
\varepsilon^\pm &: D \longrightarrow \Gamma(V_+^{0,1} \otimes V_-^{0,1}) \\
\varepsilon^- &: D \longrightarrow \Gamma(\wedge^2 V_-^{0,1})
\end{aligned}$$

be the components of ε *and for each of them let* $\varepsilon^\bullet = \sum_{k=1}^\infty \frac{1}{k!}\varepsilon_k^\bullet t^k$ *be the corresponding series expansion. Then the obstructions to finding a generalized complex structure* $\mathcal{J}_{2\varepsilon}$ *which makes* $(\mathcal{J}_{1\varepsilon}, \mathcal{J}_{2\varepsilon})$ *into a generalized Kähler structure lie in the space*

$$\mathcal{H}^{(2,1)+(1,2)} = \frac{\ker(d_{L_2} : \Gamma(\wedge^{2,1}\overline{L_2} \oplus \wedge^{1,2}\overline{L_2}))}{\mathrm{Im}\,(d_{L_2} : \Gamma(\wedge^{1,1}\overline{L_2}))},$$

and the first obstruction class is $[\overline{\partial}_+\overline{\varepsilon_1^-} + \partial_-\varepsilon_1^+]$.

Proof. According to Lemma 8.4, there is an element $a \in \Gamma(\wedge^2 L_1 \oplus \wedge^2\overline{L_1})_\mathbb{R}$ such that

$$e_*^{-a}\mathcal{J}_1 e_*^a = e_*^{-\varepsilon}\mathcal{J}_1 e^{-\varepsilon}$$

and, as in Theorem 8.3, one can compose the orthogonal transformation e_*^a with an orthogonal tranformation e_*^b which preserves $\mathcal{J}_1$ and still obtain the same deformation, that is, for any $b \in \Gamma(L_1 \otimes \overline{L_1})_\mathbb{R}$, $\mathcal{J}_{1\varepsilon}$ and $\mathcal{J}_{2\varepsilon}$ are given by

$$\mathcal{J}_{1\varepsilon} = e_*^{-a}\mathcal{J}_1 e_*^a \qquad \text{and} \qquad \mathcal{J}_{2\varepsilon} = e^{-a}e^{-b}\mathcal{J}_2 e^b e^a.$$

As before, our quest is to find b for which $\mathcal{J}_{2\varepsilon}$ is integrable and once again we do so using a power series argument. The canonical bundle of $\mathcal{J}_{2\varepsilon}$ is given by $e^a e^b U^{0,n}$ and integrability is equivalent to

$$e^{-b}e^{-a}d^H e^a e^b : \mathcal{U}^{0,n} \longrightarrow \mathcal{U}^{n-1,1} \oplus \mathcal{U}^{n-1,-1}.$$

If we decompose a and b into their $V_\pm^{1,0}$ and $V_\pm^{0,1}$ components as in (8.2) and (8.3) and write the corresponding components in power series, then the first condition we must solve is the linear vanishing of the components of $\overline{N_{\mathcal{J}_{2\varepsilon}}}$, the Nijenhuis tensor of $\mathcal{J}_{2\varepsilon}$

$$e^{-b}e^{-a}d^H e^a e^b : \mathcal{U}^{0,n} \longrightarrow \mathcal{U}^{1,n-3} \oplus \mathcal{U}^{-1,n-3}.$$

The linear term which maps $\mathcal{U}^{0,n}$ into $\mathcal{U}^{-1,n-3} \oplus \mathcal{U}^{1,n-3}$ is

$$d_{L_2}\beta_1^\pm + \partial_-\alpha_1^+ + \overline{\partial}_+\overline{\alpha_1^-}$$

So we see that the vanishing of the linear part of the Nijenhuis tensor is equivalent the following condition

$$d_{L_2}\beta_1^\pm = -(\overline{\partial}_+\overline{\alpha_1^-} + \partial_-\alpha_1^+). \tag{8.6}$$

Integrability of $\mathcal{J}_{1\varepsilon}$ means that the $\mathcal{U}^{\pm 3,n-3}$ components of $e^{-a}e^{-b}d^H e^a e^b|_{\mathcal{U}^{0,n}}$ vanish and the linear part of these components is $\overline{\partial}_+\alpha_1^+ : \mathcal{U}^{0,n} \longrightarrow \mathcal{U}^{-3,n-3}$ and $\partial_-\overline{\alpha_1^-} : \mathcal{U}^{0,n} \longrightarrow$

$\mathcal{U}^{3,n-3}$, hence α_1^+ is $\overline{\partial}_+$-closed and $\overline{\alpha_1^-}$ is ∂_--closed, showing that $\overline{\partial}_+\overline{\alpha_1^-} + \partial_-\alpha_1^+$ is d_{L_2}-closed and hence represents a class in $\mathcal{H}^{(2,1)+(1,2)}$. If this class vanishes, we can find $\beta_1^\pm$ which solves (8.6). That is the first obstruction class is

$$[\overline{\partial}_+\overline{\varepsilon_1^-} + \partial_-\varepsilon_1^+] \in \mathcal{H}^{(2,1)+(1,2)}.$$

Next we assume that we have chosen $\beta_j^\pm$ for $j < k$ such that the Nijenhuis tensor of $\mathcal{J}_2$ vanishes to order $k-1$ and we must choose $\beta_k^\pm$ which makes the Nijenhuis tensor of $\mathcal{J}_2$ vanish to order k. Expanding the operator $e^{-b}e^{-a}d^He^ae^b$ to order k we see that the degree k part of the Nijenhuis tensor is:

$$(\overline{N_{\mathcal{J}_2\varepsilon}})_k = d_{L_2}\beta_k^\pm + \partial_-\alpha_k^+ + \overline{\partial}_+\overline{\alpha_k^-} + F_k(a_1,\cdots,a_{k-1},b_1,\cdots,b_{k-1}), \tag{8.7}$$

for some function F_k which takes values in $\Gamma(\wedge^{2,1}\overline{L_2}\oplus\wedge^{1,2}\overline{L_2})$, and the vanishing of $\overline{N_{\mathcal{J}_2\varepsilon}}$ to order k is equivalent to the requirement that the term above vanishes, which can only be achived if

$$d_{L_2}\beta_k^\pm = -(\partial_-\alpha_k^+ + \overline{\partial}_+\overline{\alpha_k^-} + F_k(a_1,\cdots,a_{k-1},b_1,\cdots,b_{k-1})).$$

Next we prove that the right hand side above is d_{L_2}-closed, which allows us to conclude that the obstruction to find $\beta_k^\pm$ lies in $\mathcal{H}^{(2,1)+(1,2)}$.

We let

$$b_{<k+1} = \sum_{j<k+1}\tfrac{1}{j!}(\beta_j^\pm + \overline{\beta^\pm}_j)t^j.$$

Since $e^{-a}\mathcal{J}_1e^a = e^{-a}e^{-b_{<k+1}}\mathcal{J}_1e^{b_{<k+1}}e^a$ is integrable for any choice of b_k, we have that the operator

$$e^{-b_{<k+1}}e^{-a}d^He^ae^{b_{<k+1}}|_{\mathcal{U}^{0,n}}$$

has two components, say $d^H|_{\mathcal{U}^{n,0}} = \overline{\partial}_{\mathcal{J}_2\varepsilon} + \overline{N_{\mathcal{J}_2\varepsilon}}$,

$$\overline{\partial}_{\mathcal{J}_2k} = \overline{\partial}_{\mathcal{J}_2} + \overline{v}(t), \qquad \overline{v(t)} \in \Gamma(\overline{L_2}),\ \overline{v}(0) = 0;$$
$$\overline{N_{\mathcal{J}_2\varepsilon}} = (\overline{N_{\mathcal{J}_2\varepsilon}})_kt^k + O(t^{k+1}) \in \Gamma(\wedge^{1,2}\overline{L_2}).$$

According to Corollary 2.4, $\{\overline{\partial}_{\mathcal{J}_2\varepsilon}, \overline{N_{\mathcal{J}_2\varepsilon}}\} = 0$, hence, in particular, the degree k part of this operator vanishes, i.e.,

$$0 = \{\overline{\partial}_{\mathcal{J}_2}, (\overline{N_{\mathcal{J}_2\varepsilon}})_k\} = d_{L_2}(\overline{N_{\mathcal{J}_2\varepsilon}})_k.$$

□

Remark. There are natural maps

$$\pi_{2,1}: \mathcal{H}^{(2,1)+(1,2)} \longrightarrow H^{2,1}_{\overline{\partial}_+}(M;\overline{L_2}) \qquad \textit{and} \qquad \pi_{1,2}: \mathcal{H}^{(2,1)+(1,2)} \longrightarrow H^{1,2}_{\partial_-}(M;\overline{L_2}).$$

given by taking the appropriate components of a class in $\mathcal{H}^{(2,1)+(1,2)}$. It is possible, however, that a nontrivial class in $\mathcal{H}^{(2,1)+(1,2)}$ is mapped into the trivial class by both projections. In terms of deformations, this means that there may be no obstruction to deforming a generalized Kähler structure as either a positive or a negative SKT structure, while the generalized Kähler deformation problem itself is obstructed. Next theorem gives a concrete situation where this phenomenon happens.

Theorem 8.9. *Let* (M, ω, I) *be a compact Kähler manifold,* ω_t *be a family of deformations of the symplectic structure and* $\mathcal{J}_{1t}$ *be the corresponding family of generalized complex structures. Then, for* t *small, one can deform* I *into an generalized almost complex structure,* $\mathcal{J}_{2t}$*, such that* $(\mathcal{J}_{1t}, \mathcal{J}_{2t})$ *is a generalized complex extension of a positive SKT structure.*

Proof. Due to Moser's trick, it is enough to consider linear deformations of ω by a harmonic 2-form ε. Further if we let $\varepsilon^{p,q}$ be the components of ε according to the usual (p, q) decomposition of forms determined by the complex structure, then $\omega + t\varepsilon^{1,1}$ is still a Kähler form with respect to I, so it is enough tho prove the result for a deformation given by a harmonic form $\mathrm{Re}(\varepsilon^{2,0})$.

Next we observe that it is enough to prove the result for a deformation of the form $\omega + t\varepsilon^{0,2}$, with $\varepsilon^{0,2} \in \Omega^{0,2}(M)$ a closed form. Indeed, let $\mathcal{J}_{1t}$ be the generalized complex structure corresponding to the complex valued form $\omega + t\varepsilon^{0,2}$, that is, $\mathcal{J}_{1t}$ has canonical bundle $e^{i(\omega + t\varepsilon^{0,2})}$. If we can find a family of generalized almost complex structures $\mathcal{J}_{2t}$ which agrees with $\mathcal{J}_2$ at time zero such that $(\mathcal{J}_{1t}, \mathcal{J}_{2t})$ is a generalized complex extension of an SKT structure, then by performing a B-field transform with $B = t\mathrm{Im}\,(\varepsilon^{0,2})$ we get a generalized complex extension of an SKT structure with $\mathcal{J}_{1t}$ determined by $\omega + t\mathrm{Re}(\varepsilon^{0,2})$.

Since we have an isomorphism of Lie algebroids

$$L_1 = \{X - i\omega(X) : X \in T_{\mathbb{C}}M\} \cong T_{\mathbb{C}}M,$$

given by the projection onto $T_{\mathbb{C}}M$, we see that the differential complex $(\Gamma(\wedge^{\bullet}\overline{L_1}), d_{L_1})$ is isomorphic to $(\Omega^{\bullet}(M), d)$. In fact, the map ψ from Proposition 1.9 gives the isomorphism:

$$\psi : \wedge^{\bullet}T^*M \longrightarrow \wedge^{\bullet}\overline{L_1} \qquad \psi(\varphi) = e^{i\omega}e^{\frac{i}{2}\omega^{-1}}\varphi.$$

And according to Proposition 5.14, the operators δ_+, δ_-, $\overline{\delta}_+$ and $\overline{\delta}_-$ are related to ∂, $\overline{\partial}$, $\partial^{\omega^{-1}}$ and $\overline{\partial}^{\omega^{-1}}$ via Ψ:

$$\Psi^{-1}\delta_+\Psi = \frac{i}{2}\overline{\partial}^{\omega^{-1}}; \qquad \Psi^{-1}\overline{\delta}_+\Psi = \partial;$$

$$\Psi^{-1}\delta_-\Psi = \frac{i}{2}\partial^{\omega^{-1}}; \qquad \Psi^{-1}\overline{\delta}_-\Psi = \overline{\partial}.$$

Since $\varepsilon \in \Omega^{0,2}(M)$, then $\psi(\varepsilon) \in \Gamma(\wedge^2 L_2) \cap \Gamma(\wedge^2\overline{L_1}) = \Gamma(\wedge^2 V_-^{0,1})$ represents a d_{L_2}-cohomology class which gives rise to the deformation of $\mathcal{J}_1$ corresponding to the complex form $\omega + t\varepsilon$.

According to Proposition 8.5, there is a smooth family $\mathcal{J}_{2t}$ such that $(\mathcal{J}_{1t}, \mathcal{J}_{2t})$ is a generalized complex extension of an SKT structure. □

Remark. Of course, same result holds for negative SKT structures. That is any deformation of the symplectic structure on a Kähler manifold can be completed to a generalized Hermitian structure which is an extension of either a positive or a negative SKT structure.

Notice that, in effect, this theorem considers a Kähler structure as a bihermitian structure with $I_+ = I_-$ and the crucial step is the deformation of the symplectic structure by a harmonic form $\varepsilon \in \Omega^{2,0}(M)$. At that step, we use ε to deform I_- into a (possibly) nonintegrable complex structure still compatible with the same metric for which $\mathcal{J}_{1\varepsilon}$ is the desired symplectic structure. The SKT structure (g, I_+) is just the original Kähler structure.

References

[1] M. Atiyah and G. Segal, *Twisted K-theory and cohomology*, Inspired by S. S. Chern, Nankai Tracts Math., vol. 11, World Sci. Publ., Hackensack, NJ, 2006, pp. 5–43.

[2] J.-M. Bismut, *A local index theorem for non-Kähler manifolds*, Math. Ann. **284** (1989), no. 4, 681–699.

[3] H. Bursztyn, G. R. Cavalcanti, and M. Gualtieri, *Reduction of Courant algebroids and generalized complex structures*, Adv. Math. **211** (2007), no. 2, 726–765.

[4] H. Bursztyn, G. R. Cavalcanti, and M. Gualtieri, *Generalized Kähler geometry of instanton moduli spaces*, 2012, `http://arxiv.org/abs/1203.2385`.

[5] G. R. Cavalcanti, *New aspects of the dd^c-lemma*, Ph.D. thesis, Oxford University, 2004.

[6] G. R. Cavalcanti, *The decomposition of forms and cohomology of generalized complex manifolds*, J. Geom. Phys. **57** (2006), no. 1, 121–132.

[7] G. R. Cavalcanti, *Reduction of metric structures on Courant algebroids*, J. Symplectic Geom. **4** (2006), no. 3, 317–343.

[8] P. de Bartolomeis, *Symplectic deformations of Kähler manifolds*, J. Symplectic Geom. **3** (2005), no. 3, 341–355.

[9] B. de Wit and P. van Nieuwenhuizen, *Rigidly and locally supersymmetric two-dimensional nonlinear σ-models with torsion*, Nuclear Phys. B **312** (1989), no. 1, 58–94.

[10] G. W. Delius, M. Roček, A. Sevrin, and P. van Nieuwenhuizen, *Supersymmetric sigma models with nonvanishing Nijenhuis tensor and their operator product expansion*, Nuclear Phys. B **324** (1989), no. 2, 523–531.

[11] N. Enrietti, A. Fino, and L. Vezzoni, *Tamed Symplectic forms and SKT metrics*, 2010, `http://arxiv.org/abs/1002.3099`.

[12] A. Fino, M. Parton, and S. Salamon, *Families of strong KT structures in six dimensions*, Comment. Math. Helv. **79** (2004), no. 2, 317–340.

[13] A. Fino and A. Tomassini, *Blow-ups and resolutions of strong Kähler with torsion metrics*, Adv. Math. **221** (2009), no. 3, 914–935.

[14] S. J. Gates, Jr., C. M. Hull, and M. Roček, *Twisted multiplets and new supersymmetric nonlinear σ-models*, Nuclear Phys. B **248** (1984), no. 1, 157–186.

[15] P. Gauduchon, *Le théorème de l'excentricité nulle*, C. R. Acad. Sci. Paris Sér. A-B **285** (1977), no. 5, A387–A390.

[16] R. Goto, *Deformations of generalized complex and generalized Kähler structures*, J. Differential Geom. **84** (2010), no. 3, 525–560.

[17] J. Grabowski, *Courant-Nijenhuis tensors and generalized geometries*, Groups, geometry and physics, Monogr. Real Acad. Ci. Exact. Fís.-Quím. Nat. Zaragoza, 29, Acad. Cienc. Exact. Fís. Quím. Nat. Zaragoza, Zaragoza, 2006, pp. 101–112.

[18] G. Grantcharov, G. Papadopoulos, and Y. S. Poon, *Reduction of HKT-structures*, J. Math. Phys. **43** (2002), no. 7, 3766–3782.

[19] M. Gualtieri, *Generalized geometry and the Hodge decomposition*, 2004, `http://arxiv.org/pdf/math/0409093v1`.

[20] M. Gualtieri, *Generalized Kähler geometry*, 2010, `http://arxiv.org/pdf/1007.3485v1`.

[21] M. Gualtieri, *Generalized complex geometry*, Ann. of Math. (2) **174** (2011), no. 1, 75–123.

[22] N. Hitchin, *Brackets, forms and invariant functionals*, Asian J. Math. **10** (2006), no. 3, 541–560.

[23] C. M. Hull and E. Witten, *Supersymmetric sigma models and the heterotic string*, Phys. Lett. B **160** (1985), no. 6, 398–402.

[24] K. Kodaira and D. C. Spencer, *On deformations of complex analytic structures. III. Stability theorems for complex structures*, Ann. of Math. (2) **71** (1960), 43–76.

[25] Y. Kosmann-Schwarzbach, *Derived brackets*, Lett. Math. Phys. **69** (2004), 61–87.

[26] J.-L. Koszul, *Crochet de Schouten-Nijenhuis et cohomologie*, Astérisque (1985), no. Numero Hors Serie, 257–271. The mathematical heritage of Élie Cartan (Lyon, 1984).

[27] Z.-J. Liu, A. Weinstein, and P. Xu, *Manin triples for Lie bialgebroids*, J. Differential Geom. **45** (1997), no. 3, 547–574.

[28] M. Lübke and A. Teleman, *The Kobayashi-Hitchin correspondence*, World Scientific Publishing Co. Inc., River Edge, NJ, 1995.

[29] M. L. Michelsohn, *Clifford and spinor cohomology of Kähler manifolds*, Amer. J. Math. **102** (1980), no. 6, 1083–1146.

[30] A. Neveu and J. H. Schwarz, *Factorizable dual model of pions*, Nuclear Physics B **31** (1971), no. 1, 86 – 112.

[31] J. Wess and B. Zumino, *Consequences of anomalous Ward identities*, Phys. Lett. **37B** (1971), 95–97.